


Math 4300

Homework #4

Solutions

---



①  $A = (-1, -2), B = (2, 1), C = (0, -1)$

(a) They don't lie on a vertical line.

What about a line  $L_{m,b}$ ?

Plug them into  $y = mx + b$  to get:

$-2 = -m + b$	① ← A plugged in
$1 = 2m + b$	② ← B plugged in
$-1 = b$	③ ← C plugged in



$b = -1$  gives then  $m = 1$  in both ① and ②.


Let's verify that all three points satisfy the equation  $y = x - 1$ .

We have:

$$\begin{aligned} -2 &= -1 - 1 && \checkmark \\ 1 &= 2 - 1 && \checkmark \\ -1 &= 0 - 1 && \checkmark \end{aligned}$$

These three points all lie on  $L_{m,b} = L_{1,-1}$ .

So they are collinear.



### Method 1 - by def

(b) In the picture on the previous page we can guess that  $A-C-B$  is true.

Let's check:

(i) we have three distinct points ✓

(ii)  $A, B, C$  are collinear ✓

(iii)  $d_E(A, C) + d_E(C, B)$

$$= \sqrt{(-1-0)^2 + (-2+1)^2} + \sqrt{(0-2)^2 + (-1-1)^2}$$

$$= \sqrt{2} + \sqrt{8} = \sqrt{2}(1+2) = 3\sqrt{2}$$

$$A = (-1, -2)$$

$$B = (2, 1)$$

$$C = (0, -1)$$

$$\begin{aligned}\text{And, } d_E(A, B) &= \sqrt{(-1-2)^2 + (-2-1)^2} \\ &= \sqrt{9+9} \\ &= \sqrt{18} = 3\sqrt{2}\end{aligned}$$

$$\text{So, } d_E(A, C) + d_E(C, B) = d_E(A, B).$$

By conditions (i), (ii), (iii) we have that  $A-C-B$  is true.

By a later problem of this HW we cannot have also  $A-B-C$  or  $B-A-C$ . Thus, only  $A-C-B$  is true.

### method 2

Note: There is another way to check condition (iii) above. By using the standard ruler!

The standard ruler on  $L_{1,-1}$ .

The standard ruler is  $f: L_{1,-1} \rightarrow \mathbb{R}$

$$\text{where } f(x, y) = x\sqrt{1+1^2} = \sqrt{2}x$$




Apply  $f$  to  $A, B, C$  to get

$$f(A) = f(-1, -2) = -\sqrt{2}$$

$$f(B) = f(2, 1) = 2\sqrt{2}$$

$$f(C) = f(0, -1) = 0\sqrt{2} = 0$$



Since  $f(A) < f(C) < f(B)$  we have  
 $A - C - B$  by a theorem from class.

(c) Since  $A - C - B$ , then by a  
theorem in class we also  
have  $B - C - A$ .

②  $A = (1, 2), B = (3, 4), C = (4, \sqrt{19})$

(a) They aren't on a vertical line.

Let's see if they are on some  $C_r$ .

Plug  $A, B$  into  $(x-c)^2 + y^2 = r^2$  to get:

(we only need to first find  $\overleftrightarrow{AB}$  and then check if  $C$  lies on it also)

$$\begin{aligned} (1-c)^2 + 2^2 &= r^2 \\ (3-c)^2 + 4^2 &= r^2 \end{aligned}$$

①  $\leftarrow$  A plugged in

②  $\leftarrow$  B plugged in



$$\begin{aligned} c^2 - 2c + 5 &= r^2 \\ c^2 - 6c + 25 &= r^2 \end{aligned}$$

①

②

① - ② gives  $4c - 20 = 0$ . So,  $c = 5$ .

Plugging  $c = 5$  into ① gives  $r = \sqrt{20}$

Now let's see if all three points

satisfy  $(x-5)^2 + y^2 = 20$

$$(1-5)^2 + 2^2 = 20 \quad \checkmark$$

$$(3-5)^2 + 4^2 = 20 \quad \checkmark$$

$$(4-5)^2 + \sqrt{19}^2 = 20 \quad \checkmark$$


$$A = (1, 2)$$

$$B = (3, 4)$$

$$C = (4, \sqrt{19})$$

So, A, B, C all lie on  $5 \sqrt{20}$

Note  $\sqrt{20} \approx 4.47$  and  $\sqrt{19} \approx 4.36$



Method 1 - by def

(b) From the picture it looks like

A-B-C. Let's verify.

(i) we have three distinct points

(ii) A, B, C are collinear

$$(iii) d_H(A, B) + d_H(B, C)$$

$$\begin{aligned} A &= (1, 2) \\ B &= (3, 4) \\ C &= (4, \sqrt{19}) \end{aligned}$$

recall: If

$$P = (x_1, y_1)$$

$$Q = (x_2, y_2)$$

are on  $c^L_r$

$$d(P, Q) = \left| \ln \left( \frac{(x_1 - c + r)/y_1}{(x_2 - c + r)/y_2} \right) \right|$$

$$= \left| \ln \left( \frac{(1-5+\sqrt{20})/2}{(3-5+\sqrt{20})/4} \right) \right| + \left| \ln \left( \frac{(3-5+\sqrt{20})/4}{(4-5+\sqrt{20})/\sqrt{19}} \right) \right|$$

$$= \left| \ln \left( \frac{(-4+\sqrt{20})/2}{(-2+\sqrt{20})/4} \right) \right| + \left| \ln \left( \frac{(-2+\sqrt{20})/4}{(-1+\sqrt{20})/\sqrt{19}} \right) \right|$$

$$\approx -2.34 < 0$$

$$\approx -0.60 < 0$$

$$= -\ln \left( \frac{(-4+\sqrt{20})/2}{(-2+\sqrt{20})/4} \right) - \ln \left( \frac{(-2+\sqrt{20})/4}{(-1+\sqrt{20})/\sqrt{19}} \right)$$

the negative signs are coming from dropping the abs value since the # inside the abs value is negative

$$= \ln \left( \frac{(-2+\sqrt{20})/4}{(-4+\sqrt{20})/2} \right) + \ln \left( \frac{(-1+\sqrt{20})/\sqrt{19}}{(-2+\sqrt{20})/4} \right)$$

$$-\ln(c) = \ln\left(\frac{1}{c}\right)$$

$$\ln(A) + \ln(B) = \ln(AB)$$

$$= \ln \left( \frac{(-1+\sqrt{20})/\sqrt{19}}{(-4+\sqrt{20})/2} \right)$$

$$\approx 0.963 > 0$$

$$= \left| \ln \left( \frac{(-1+\sqrt{20})/\sqrt{19}}{(-4+\sqrt{20})/2} \right) \right|$$

$$= \left| \ln \left( \frac{(4-5+\sqrt{20})/19}{(1-5+\sqrt{20})/2} \right) \right|$$

$$= d_H(C, A)$$

$$= d_H(A, C)$$

property of distance functions

Thus, from (i), (ii), (iii) we see that  
 $A-B-C$ . From a hw problem  
is this topic we can't also have  
 $A-C-B$  or  $B-A-C$ .

**Method 2** - There is another way to check  
condition (iii). Use the standard ruler!  
The standard ruler on  ${}_5L_{\sqrt{20}}$  is given by  
 $f: {}_5L_{\sqrt{20}} \rightarrow \mathbb{R}$  where  $f(x, y) = \ln\left(\frac{x-5+\sqrt{20}}{y}\right)$

We have

$$f(A) = f(1, 2) = \ln\left(\frac{1-5+\sqrt{20}}{2}\right) \approx -1.44$$


$$f(B) = f(3, 4) = \ln\left(\frac{3-5+\sqrt{20}}{4}\right) \approx -0.48$$

$$f(C) = f(4, \sqrt{19}) = \ln\left(\frac{4-5+\sqrt{20}}{\sqrt{19}}\right) \approx -0.227$$


Since  $f(C) < f(B) < f(A)$

We know  $C-B-A$  or equivalently  
from a theorem from class

We have  $A-B-C$ .



③ (a) All three points  $A=(1,2)$ ,  $B=(1,4)$ , and  $C=(1,5)$  lie on  $L$  and so they are collinear.



(b) Let's check if  $A-B-C$ ,  $A-C-B$ , or  $B-A-C$ .

You can use the def way of doing this like in the previous problems but this time let's use the easier standard ruler method (method 2 in the previous problems)

(i)  $A, B, C$  are distinct points ✓

(ii)  $A, B, C$  are collinear ✓

(iii) the standard ruler on  $L$  is given by  $f(l, y) = \ln(y)$

We have


$$f(A) = f(l, 2) = \ln(2) \approx 0.693$$

$$f(B) = f(l, 4) = \ln(4) \approx 1.386$$

$$f(C) = f(l, 5) = \ln(5) \approx 1.609$$

Since  $f(A) < f(B) < f(C)$

we know  $A - B - C$ .





④ Let  $A, B$  be points with  $A \neq B$  in a metric geometry.

Let  $C \in \overleftrightarrow{AB}$ .

Let  $f: \overleftrightarrow{AB} \rightarrow \mathbb{R}$  be a ruler for  $\overleftrightarrow{AB}$ .

Since  $A \neq B$  and  $f$  is a bijection we know that  $f(A) \neq f(B)$ .

Then there are several cases to consider:

$$(i) f(A) < f(B) < f(C)$$

$$(ii) f(A) < f(C) < f(B)$$

$$(iii) f(A) = f(C)$$

$$(iv) f(B) < f(A) < f(C)$$

$$(v) f(B) < f(C) < f(A)$$

$$(vi) f(B) = f(C)$$

$$(vii) f(C) < f(A) < f(B)$$

$$(viii) f(C) < f(B) < f(A)$$

If (i) is true then  $A = B = C$ .

If (ii) is true then  $A = C = B$

If (iii) is true then  $A = C$ .

If (iv) is true then  $B = A = C$

which implies that  $C = A = B$ .

If (v) is true then  $B = C = A$

which implies that  $A = C = B$ .

If (vi) is true then  $B = C$

If (vii) is true then  $C = A = B$ .

If (viii) is true then  $C = B = A$ ,

which implies that  $A = B = C$ .

Thus summarizing either:

$C = A = B$ , or  $C = A$ , or  $A = C = B$ ,

or  $C = B$ , or  $A = B = C$ .

And no two of these can be true at the same time (by the above cases)

since  $f$  is 1-1

from a thm in class:  
If  $x = y = z$  then  $z = y = x$




⑤ Let  $l$  be a line and  $A, B, C$  be distinct points on  $l$  in a metric geometry.

So,  $A \neq B$ ,  $A \neq C$ , and  $B \neq C$ .

We know  $l = \overleftrightarrow{AB}$  since there is a unique line through any two distinct points.

By the previous HW problem we know that there are only three possible outcomes (since  $C \neq A$  and  $C \neq B$ )

These are that either  $C-A-B$ , or  $A-C-B$ , or  $A-B-C$ .

Since  $C-A-B$  implies  $B-A-C$ , this gives that the only three possibilities are  $B-A-C$ , or  $A-C-B$ , or  $A-B-C$ . 

⑥

Suppose that  $A-B-C$  and  $B-C-D$  in some metric geometry.

Since  $A-B-C$  we know  $A, B, C$  are distinct and collinear and all lie on  $\overleftrightarrow{BC}$ .

Since  $B-C-D$  we know that  $B, C, D$  are all distinct and collinear and all lie on  $\overleftrightarrow{BC}$ .

Let  $f: \overleftrightarrow{BC} \rightarrow \mathbb{R}$  be a ruler.

Since  $A-B-C$  we know either

$$(i) f(A) < f(B) < f(C)$$

$$\text{or } (ii) f(C) < f(B) < f(A).$$

Since  $B-C-D$  we know either

$$(iii) f(B) < f(C) < f(D)$$

$$\text{or } (iv) f(D) < f(C) < f(B).$$

Case 1:

Suppose (i)  $f(A) < f(B) < f(C)$  is true.

Then, we can't have (iv) since then  $f(C) < f(B)$ .

So we must have (iii), that is  $f(B) < f(C) < f(D)$ .

Thus,  $f(A) < f(B) < f(C) < f(D)$ .

So,  $A-B-D$  and  $A-C-D$ .

Case 2:

Suppose (ii)  $f(C) < f(B) < f(A)$  is true.

Then, we can't have (iii) since then  $f(B) < f(C)$ .

So we must have (iv), that is  $f(D) < f(C) < f(A)$ .

Thus,  $f(D) < f(C) < f(B) < f(A)$ .

Hence,  $D-B-A$  and  $D-C-A$ .

So,  $A-B-D$  and  $A-C-D$ .



⑦ Suppose  $A-C-D$  and  $A-C-B$  is some metric geometry.

Then,  $A, B, C, D \in \overleftrightarrow{AC}$ .

Let  $l = \overleftrightarrow{AC}$  and  $f: l \rightarrow \mathbb{R}$  be a ruler.

Since  $A-C-D$  we get either

$$(i) f(A) < f(C) < f(D)$$

$$\text{or } (ii) f(D) < f(C) < f(A)$$

Case (i): Suppose  $f(A) < f(C) < f(D)$ .

Since  $A-C-B$  we have either

$$f(A) < f(C) < f(B) \quad (*)$$

$$\text{or } f(B) < f(C) < f(A) \quad (**)$$

But  $(**)$  can't happen since we are assuming that  $f(A) < f(C)$ .

Thus we must have  $(*)$ .

Combining the case (i) conditions with  $(*)$

We get either

$$f(A) < f(C) < f(D) < f(B)$$

$$\text{or } f(A) < f(C) < f(B) < f(D).$$

In the first inequality we get  $A-D-B$ .

In the second we get  $A-B-D$ .

So either  $A-D-B$  or  $A-B-D$ .

Case (ii): Suppose  $f(D) < f(C) < f(A)$ .

Since  $A-C-B$  we have either

$$f(A) < f(C) < f(B)$$

(\*\*\*)

$$\text{or } f(B) < f(C) < f(A)$$

(\*\*\*\*)

We see that (\*\*\*) can't happen because we are assuming that  $f(C) < f(A)$ .

Thus we must have (\*\*\*\*)

Combining case (ii) with (\*\*\*\*) we get


that either

$$f(D) < f(B) < f(C) < f(A)$$

$$\text{or } f(B) < f(D) < f(C) < f(A).$$

Thus either  $D-B-A$  or  $B-D-A$ .

So either  $A-B-D$  or  $A-D-B$ .





⑧ Suppose  $A-D-C$  and  $A-C-B$ .

Then  $A, B, C, D$  all lie on  $l = \overleftrightarrow{AC}$ .

Let  $f: l \rightarrow \mathbb{R}$  be a ruler.

Since  $A-D-C$  we have either

$$(i) \quad f(A) < f(D) < f(C)$$

$$\text{or } (ii) \quad f(C) < f(D) < f(A)$$

Case (i): Suppose  $f(A) < f(D) < f(C)$ .

Since  $A-C-B$  we have either

$$f(A) < f(C) < f(B) \quad (*)$$

$$\text{or } f(B) < f(C) < f(A) \quad (**)$$

We can't have  $(**)$  since we are assuming in this case that  $f(A) < f(C)$ .

Thus we have  $(*)$ .

We get that case (i) and (\*) give  
that  $f(A) < f(D) < f(C) < f(B)$ .

Thus,  $A - D - B$ .

Case (ii): Suppose  $f(C) < f(D) < f(A)$ .

Since  $A - C - B$  we have either

$$f(A) < f(C) < f(B) \quad (***)$$

$$\text{or } f(B) < f(C) < f(A) \quad (****)$$

We can't have (\*\*\*) since we are assuming  
in this case that  $f(C) < f(A)$ .

Thus we have (\*\*\*\*).

We get that case (ii) and (\*\*\*\*) give

$$\text{that } f(B) < f(C) < f(D) < f(A).$$

Thus,  $B - D - A$ .

So,  $A - D - B$ .

In either case we get  $A - D - B$ . 

(9) Suppose that  $A-Q-B$ ,  $A-P-B$ ,  
and  $P-C-Q$ . Let  $l = \overleftrightarrow{AB}$ .

Since  $A-Q-B$  and  $A-P-B$  we know  
all of  $A, B, P, Q$  lie on  $l$ .

Let  $f: l \rightarrow \mathbb{R}$  be a ruler for  $l$ .

Since  $A-Q-B$  we get two cases:

either  $f(A) < f(Q) < f(B)$  or  $f(B) < f(Q) < f(A)$ .

I'll prove this problem for when

$f(A) < f(Q) < f(B)$ , you try the other case.

Suppose  $f(A) < f(Q) < f(B)$ . (\*)

Since  $A-P-B$  we have two cases.

case 1: Suppose  $f(A) < f(P) < f(B)$ . (1)

Since  $P-C-Q$  we have either

$f(P) < f(C) < f(Q)$  or  $f(Q) < f(C) < f(P)$ .  
(i) (ii)

If (i), then

$f(A) < f(P) < f(C) < f(Q) < f(B)$   
(1) (i) (ii) (\*)

So,  $A-C-B$ .

If (ii), then

$$f(A) < f(Q) < f(C) < f(P) < f(B)$$

(\*)
(ii)
(iii)
(i)

So, A-C-B.

Case 2: Suppose  $f(B) < f(P) < f(A)$  (2)

Since  $P-C-Q$  we have either  
 $\underbrace{f(P) < f(C) < f(Q)}_{(i)}$  or  $\underbrace{f(Q) < f(C) < f(P)}_{(ii)}$ .

If (i), then

$$f(B) < \underset{(2)}{f(P)} < \underset{(1)}{f(C)} < \underset{(1)}{f(Q)} < \underset{(*)}{f(B)}$$

This is a contradiction, so this case can't happen.

If (ii), then


(ii), then

$$f(A) < f(Q) < f(B) < f(P) < f(A)$$

(\*)                      (\*)                      (2)                      (2)

Hence case

This is a contradiction, so this case can't happen.




(10) Let  $A, B, C \in \mathbb{R}^2$  be distinct points.

( $\Rightarrow$ ) Suppose  $A-B-C$ .

Then,  $A, B, C$  all lie on  $l = \overleftrightarrow{AB} = \overleftrightarrow{AC} = \overleftrightarrow{BC}$ .

Since  $B \in l = \overleftrightarrow{AC}$  and  $\overleftrightarrow{AC} = L_{AC}$  we



know that  $B = A + t(C - A)$  where  $t \in \mathbb{R}$ .

Goal:

We must show that  $0 < t < 1$

If we can do this then we have proven the ( $\Rightarrow$ ) direction of this proof

Let  $A = (x_a, y_a)$ ,  $B = (x_b, y_b)$

and  $C = (x_c, y_c)$ .

Then,  $B = A + t(C - A)$  becomes

(\*)

$$\underbrace{(x_b, y_b)}_B = \underbrace{(x_a + tx_c - tx_a, y_a + ty_c - ty_a)}_{A + t(C - A)}$$

We now break the proof into two cases:  
if  $l$  is a vertical line and if  
 $l$  is a non-vertical line.

**Case 1:** Suppose  $l = L_d$  is a vertical line.

Let  $f: l \rightarrow \mathbb{R}$  be the standard  
ruler given by  $f(x, y) = y$ .

Apply the ruler  $f$  to  $(*)$  above to get

$$y_b = y_a + t y_c - t y_a.$$

So,

$$y_b - y_a = t(y_c - y_a) \quad (**)$$



Since  $A-B-C$  we know that either

$$(i) f(A) < f(B) < f(C)$$


or (ii)  $f(C) < f(B) < f(A)$

vertical line pic

Thus either

$$(i) y_a < y_b < y_c$$

or (ii)  $y_c < y_b < y_a$



case (i): Suppose we have (i).

That is, suppose  $y_a < y_b < y_c$ .

Then,  $0 < y_b - y_a$  and  $0 < y_c - y_a$ .

In (\*\*) we have  $\underbrace{y_b - y_a}_{>0} = t \underbrace{(y_c - y_a)}_{>0}$

So we must have  $t > 0$ .

Why can't  $t \geq 1$ ?

Suppose  $t \geq 1$ .

Then from (\*) we get

$$y_b - y_a = t(y_c - y_a) \geq y_c - y_a$$

But then  $y_b \geq y_c$ .

This contradicts  $y_b < y_c$  from (i).

Therefore  $0 < t < 1$  and we are done with this case

case (ii): Suppose (ii), that is,  $y_c < y_b < y_a$ .

Then,  $0 < y_a - y_c$  and  $0 < y_a - y_b$ .

So,  $y_c - y_a < 0$  and  $y_b - y_a < 0$ .

Thus we have from (\*)  $\underbrace{(y_b - y_a)}_{< 0} = t \underbrace{(y_c - y_a)}_{< 0}$

So,  $t > 0$ .

Why can't we have  $t \geq 1$ ?

Suppose  $t \geq 1$ .



Then  $\frac{1}{t} \leq 1$  and  $\frac{1}{t}(y_b - y_a) = (y_c - y_a)$

Thus,  $\frac{1}{t}(y_a - y_b) = y_a - y_c$ .

Then,  $y_a - y_c = \underbrace{\frac{1}{t}}_{\leq 1}(y_a - y_b) \leq y_a - y_b$ .

So,  $-y_c \leq -y_b$

Then  $y_c \geq y_b$ .

But this contradicts  $y_c < y_b$  from (ii)


Therefore,  $t \geq 1$  can't be true  
and thus  $0 < t < 1$ .

This completes the proof of case 1.

**case 2:** Suppose  $l = L_{m,c}$  is a non-vertical line.

Let  $f: l \rightarrow \mathbb{R}$  be the standard ruler.

Then,  $f(x, y) = x \sqrt{1+m^2} = xM$   
where  $M = \sqrt{1+m^2}$



Recall that (\*) says that

$$(x_b, y_b) = (x_a + t x_c - t x_a, y_a + t y_c - t y_a)$$

Applying  $f$  to the above eqn gives:

$$Mx_b = M(x_a + t x_c - t x_a)$$

Cancelling  $M$  and subtracting  $x_a$  gives


$$x_b - x_a = t(x_c - x_a) \quad (***)$$

picture

Since  $A-B-C$  we know that either

$$(i) f(A) < f(B) < f(C)$$

$$\text{or } (ii) f(C) < f(B) < f(A)$$



That is, either

$$(i) Mx_a < Mx_b < Mx_c$$

$$\text{or } (ii) Mx_c < Mx_b < Mx_a$$

Since  $M > 0$  this becomes either

$$(i) x_a < x_b < x_c$$

$$\text{or } (ii) x_c < x_b < x_a$$

case (i): Suppose we have (i).

Then  $x_b - x_a > 0$  and  $x_c - x_a > 0$ .

Thus, from (\*\*\*)  $\underbrace{x_b - x_a}_{>0} = t \underbrace{(x_c - x_a)}_{>0}$

we must have  $t > 0$ .

We can't have  $t \geq 1$  because if we did then we would get

$$x_b - x_a = t(x_c - x_a) \geq x_c - x_a$$

which would give  $x_b \geq x_c$  which isn't true by (i).

Thus,  $0 < t < 1$ . So we are done with this case.

case (ii): Suppose we have case (ii)

Then,  $x_a - x_c > 0$  and  $x_a - x_b > 0$ .

So,  $x_c - x_a < 0$  and  $x_b - x_a < 0$ .

Thus, from (\*\*\*)

$$\underbrace{x_b - x_a}_{<0} = t \underbrace{(x_c - x_a)}_{<0},$$

we must have  $t > 0$ .

Let's show  $t < 1$ .

Suppose  $t \geq 1$ .

Then,  $\frac{1}{t} \leq 1$  and  $\frac{1}{t}(x_b - x_a) = (x_c - x_a)$

So,  $\frac{1}{t}(x_a - x_b) = x_a - x_c$ . ←  $x(-1)$

Then,  $x_a - x_c = \underbrace{\frac{1}{t}}_{\leq 1} (x_a - x_b) \leq x_a - x_b$

So,  $-x_c \leq -x_b$ .

And then  $x_c \geq x_b$ .

This contradicts  $x_c < x_b$  from case (ii)

Thus,  $t < 1$  and so  $0 < t < 1$  and we are done with this case.

This concludes the proof of case 2.

Thus we have proven ( $\Rightarrow$ ) since there are only two cases.

( $\Leftarrow$ ) Let  $A, B, C$  be distinct points where  $B = A + t(C - A)$  with  $0 < t < 1$ .

This implies that  $B \in L_{AC} = \overleftrightarrow{AC}$ .

So,  $A, B, C$  are collinear.

Let  $l = L_{AC}$ .

Goal: We must show that  $A - B - C$ .

Let  $A = (x_a, y_a)$ ,  $B = (x_b, y_b)$ , and  $C = (x_c, y_c)$ .

Then  $B = A + t(C - A)$  becomes

$$\underbrace{(x_b, y_b)}_B = \underbrace{(x_a + t x_c - t x_a, y_a + t y_c - t y_a)}_{A + t(C - A)} \quad (\Delta)$$

We now break the proof into two cases:  
when  $l$  is vertical and when  
 $l$  is non-vertical.

case i: Suppose  $l = L_d$  is a vertical line

Recall that the standard ruler on  $l$  is

$$f: l \rightarrow \mathbb{R} \text{ given by } f(x, y) = y.$$

Apply  $f$  to  $(\Delta)$  above we get

$$y_b = y_a + t y_c - t y_a.$$

( $\Delta\Delta$ )

So,  $(y_b - y_a) = t(y_c - y_a)$  where  $0 < t < 1$ .

Since  $A \neq B$  and both points lie on the vertical line  $l$  we know  $y_b \neq y_a$ .

Thus, either  $y_b - y_a < 0$  or  $y_b - y_a > 0$ .

case ( $\bar{\lambda}$ ) Suppose  $y_b - y_a < 0$

Then from ( $\Delta\Delta$ ) since we have

$$\underbrace{(y_b - y_a)}_{< 0} = \underbrace{t}_{> 0} (y_c - y_a)$$

we must have  $y_c - y_a < 0$ .

Thus, since  $y_b - y_a < 0$  and  $y_c - y_a < 0$   
we have  $y_b < y_a$  and  $y_c < y_a$ .

We want to show that  $y_c < y_b$ .

Suppose instead that  $y_b \leq y_c$ .

Since  $0 < t < 1$  we know  $1 < \frac{1}{t}$ .

So, then  $\frac{1}{t}(y_b - y_a) = y_c - y_a$  from above

and we get  $\frac{1}{t}(y_a - y_b) = y_a - y_c$ . (by mult. by -1)

Then,

$$y_a - y_c = \frac{1}{t}(y_a - y_b) > y_a - y_b \geq y_a - y_c$$

$y_b \leq y_c$   
implies  
 $-y_b \geq -y_c$

But then  $y_a - y_c > y_a - y_c$ , which is a contradiction.

Thus we must have  $y_c < y_b$ .

Summarizing the above we get  $y_c < y_b < y_a$ .

Thus,  $f(c) < f(b) < f(a)$  and so  $C-B-A$ .

Since  $C-B-A$  we have  $A-B-C$ .

Case (ii): Suppose  $y_b - y_a > 0$ .

Then from  $(\Delta\Delta)$  since we have

$$\underbrace{(y_b - y_a)}_{>0} = \underbrace{t}_{>0} (y_c - y_a)$$

we must have  $y_c - y_a > 0$

Thus,  $y_b - y_a > 0$  and  $y_c - y_a > 0$ .

So,  $y_b > y_a$  and  $y_c > y_a$

We want to show that  $y_c > y_b$ .

Suppose otherwise that  $y_c \leq y_b$ .

Then,

$$y_b - y_a = t(y_c - y_a) \leq t(y_b - y_a)$$

$$< (y_b - y_a)$$

since  
 $y_c - y_a \leq y_b - y_a$   
and  $t > 0$

since  $t < 1$

This gives  $y_b - y_a < y_b - y_a$  which is a contradiction.

Thus,  $y_c > y_b$ .

So we get  $y_a < y_b < y_c$ .

So,  $f(A) < f(B) < f(C)$

Thus  $A-B-C$ . This concludes case I.



case 2: Suppose  $l = L_{c,m}$  is a non-vertical line.  
Let  $f: l \rightarrow \mathbb{R}$  be the standard ruler  
where  $f(x,y) = Mx$  where  $M = \sqrt{1+m^2} > 0$ .

Recall that  $(\Delta)$  says that

$$(x_b, y_b) = (x_a + t x_c - t x_a, y_a + t y_c - t y_a)$$

Applying  $f$  to  $(\Delta)$  gives

$$M x_b = M (x_a + t x_c - t x_a).$$

Cancelling by  $M$  and subtracting  $x_a$  gives

$$x_b - x_a = t(x_c - x_a). \quad (\Delta\Delta\Delta)$$

Since  $A$  and  $B$  are distinct and they  
lie on a non-vertical line  
we know  $x_b \neq x_a$  and so  $x_b - x_a \neq 0$ .

We break the proof into two cases:

$$x_b - x_a < 0 \quad \text{and} \quad x_b - x_a > 0.$$

case (i) Suppose  $x_b - x_a < 0$

Then from  $(\Delta\Delta\Delta)$  since we have

$$\underbrace{(x_b - x_a)}_{< 0} = \underbrace{t}_{> 0} (x_c - x_a)$$

we must have  $x_c - x_a < 0$ .

Thus, since  $x_b - x_a < 0$  and  $x_c - x_a < 0$   
we have  $x_b < x_a$  and  $x_c < x_a$ .

We want to show that  $x_c < x_b$ .

Suppose instead that  $x_b \leq x_c$ .

Since  $0 < t < 1$  we know  $1 < \frac{1}{t}$ .

So, then  $\frac{1}{t}(x_b - x_a) = x_c - x_a$  from above

and we get  $\frac{1}{t}(x_a - x_b) = x_a - x_c$ . (by mult. by -1)

Then,

$$x_a - x_c = \frac{1}{t}(x_a - x_b) > x_a - x_b \geq x_a - x_c$$

$x_b \leq x_c$   
implies  
 $-x_b \geq -x_c$

But then  $X_a - X_c > X_a - X_c$ , which is a contradiction.

Thus we must have  $X_c < X_b$ .

Summarizing the above we get  $X_c < X_b < X_a$ .

Since  $M = \sqrt{1+m^2} > 0$  we get  $MX_c < MX_b < MX_a$ .

Thus,  $f(c) < f(b) < f(a)$  and so  $C-B-A$ .

Since  $C-B-A$  we have  $A-B-C$ .

Case (ii): Suppose  $X_b - X_a > 0$ .

Then from  $(\Delta \Delta \Delta)$  since we have

$$\underbrace{(X_b - X_a)}_{>0} = \underbrace{k}_{>0} (X_c - X_a)$$

we must have  $X_c - X_a > 0$

Thus,  $X_b - X_a > 0$  and  $X_c - X_a > 0$ .

So,  $X_b > X_a$  and  $X_c > X_a$

We want to show that  $X_c > X_b$ .

Suppose otherwise that  $X_c \leq X_b$ .

since  $x_c - x_a \leq x_b - x_a$   
and  $t > 0$

Then,

$$x_b - x_a = t(x_c - x_a) \leq t(x_b - x_a)$$

$$< (x_b - x_a)$$

since  $t < 1$

This gives  $x_b - x_a < x_b - x_a$  which is a contradiction.

Thus,  $x_c > x_b$ .

So we get  $x_a < x_b < x_c$ .

Since  $M = \sqrt{1+m^2} > 0$  this gives  $Mx_a < Mx_b < Mx_c$

So,  $f(A) < f(B) < f(C)$

Thus  $A-B-C$ .

This concludes case 2.

Thus, by cases 1 and 2 we always  
get  $A-B-C$ .

So we have proven ( $\Leftarrow$ ).

