

Math 4550  
Homework 6  
Solutions


---

---

---

---

---



①(a)  $\mathbb{Z}_{12} = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}, \bar{5}, \bar{6}, \bar{7}, \bar{8}, \bar{9}, \bar{10}, \bar{11}\}$

$H = \langle \bar{4} \rangle = \{\bar{0}, \bar{4}, \bar{8}\}$

left cosets

$\bar{0} + H = \{\bar{0}, \bar{4}, \bar{8}\}$   
 $\bar{1} + H = \{\bar{1}, \bar{5}, \bar{9}\}$   
 $\bar{2} + H = \{\bar{2}, \bar{6}, \bar{10}\}$   
 $\bar{3} + H = \{\bar{3}, \bar{7}, \bar{11}\}$

right cosets

$H + \bar{0} = \{\bar{0}, \bar{4}, \bar{8}\}$   
 $H + \bar{1} = \{\bar{1}, \bar{5}, \bar{9}\}$   
 $H + \bar{2} = \{\bar{2}, \bar{6}, \bar{10}\}$   
 $H + \bar{3} = \{\bar{3}, \bar{7}, \bar{11}\}$

The left and right cosets are the same.  
The subgroup  $H$  is normal.

$\bar{0} + H = H + \bar{0}$	$\bar{1} + H = H + \bar{1}$	$\bar{2} + H = H + \bar{2}$	$\bar{3} + H = H + \bar{3}$
$\bar{0}.$	$\bar{1}.$	$\bar{2}.$	$\bar{3}.$
$\bar{4}.$	$\bar{5}.$	$\bar{6}.$	$\bar{7}.$
$\bar{8}.$	$\bar{9}.$	$\bar{10}.$	$\bar{11}.$

G

$$\textcircled{1}(b) \quad D_8 = \{1, r, r^2, r^3, s, sr, sr^2, sr^3\}$$

$$H = \langle r^2 \rangle = \{1, r^2\}$$

Recall:

$$r^4 = 1, s^2 = 1$$

$$r^k s = s r^{-k} = s r^{4-k}$$

left cosets

$$H = \{1, r^2\}$$

$$rH = \{r, r^3\}$$

$$sH = \{s, sr^2\}$$

$$srH = \{sr, sr^3\}$$

right cosets

$$H = \{1, r^2\}$$

$$Hr = \{r, r^3\}$$

$$Hs = \{s, r^2 s\} = \{s, s r^{-2}\} = \{s, sr^2\}$$

$$Hsr = \{sr, r^2 sr\} = \{sr, s r^{-2} r\} = \{sr, sr^{-1}\} = \{sr, sr^3\}$$

The left and right cosets are the same.  
Thus  $H$  is a normal subgroup of  $D_8$ .

$H$	$rH = Hr$	$sH = Hs$	$srH = Hsr$
1 •	$r$ •	$s$ •	$sr$ •
$r^2$ •	$r^3$ •	$sr^2$ •	$sr^3$ •

$D_8$

①(c)

$$D_8 = \{1, r, r^2, r^3, s, sr, sr^2, sr^3\}$$

$$H = \langle s \rangle = \{1, s\}$$

Recall:

$$r^4 = 1, s^2 = 1$$

$$r^k s = s r^{-k} = s r^{4-k}$$

left cosets

$$H = \{1, s\}$$

$$rH = \{r, rs\} = \{r, sr^{-1}\} = \{r, sr^3\}$$

$$r^2H = \{r^2, r^2s\} = \{r^2, sr^{-2}\} = \{r^2, sr^2\}$$

$$r^3H = \{r^3, r^3s\} = \{r^3, sr^{-3}\} = \{r^3, sr\}$$

right cosets

$$H = \{1, s\}$$

$$Hr = \{r, sr\}$$

$$Hr^2 = \{r^2, sr^2\}$$

$$Hr^3 = \{r^3, sr^3\}$$

The left and right cosets differ so  $H$  is not normal.

Let's see the two partitions created by drawing them.

$H$	$rH$	$r^2H$	$r^3H$
1.	r.	$r^2$ .	$r^3$ .
s.	$sr^3$ .	$sr^2$ .	sr.

partition  
of  $D_8$   
by the  
left  
cosets  
of  $H$

H	$Hr$	$Hr^2$	$Hr^3$
1.	$r.$	$r^2.$	$r^3.$
s.	$sr.$	$sr^2.$	$sr^3.$

partition  
of  $D_8$   
by the  
right  
cosets  
of  $H$

(2)

(a) From problem 1 we got that

$$\mathbb{Z}_{12}/H = \{ \bar{0}+H, \bar{1}+H, \bar{2}+H, \bar{3}+H \}$$

Where

$$\begin{aligned}\bar{0}+H &= \{ \bar{0}, \bar{4}, \bar{8} \} = \bar{4}+H = \bar{8}+H \\ \bar{1}+H &= \{ \bar{1}, \bar{5}, \bar{9} \} = \bar{5}+H = \bar{9}+H \\ \bar{2}+H &= \{ \bar{2}, \bar{6}, \bar{10} \} = \bar{6}+H = \bar{10}+H \\ \bar{3}+H &= \{ \bar{3}, \bar{7}, \bar{11} \} = \bar{7}+H = \bar{11}+H\end{aligned}$$

The identity element is  $\bar{0}+H$ .

(b)

$$(\bar{2}+H) + (\bar{3}+H) = \bar{5}+H = \bar{1}+H$$

Since

$$(\bar{1}+H) + (\bar{3}+H) = \bar{4}+H = \bar{0}+H \leftarrow \bar{0}+H \text{ is the identity element}$$

the inverse of  $\bar{3}+H$  is  $\bar{1}+H$ .

(c)

order of  $\bar{1}+H$

$$\bar{1}+H \neq \bar{0}+H$$

$$(\bar{1}+H) + (\bar{1}+H) = \bar{2}+H \neq \bar{0}+H$$

$$(\bar{1}+H) + (\bar{1}+H) + (\bar{1}+H) = \bar{3}+H \neq \bar{0}+H$$

$$(\bar{1}+H) + (\bar{1}+H) + (\bar{1}+H) + (\bar{1}+H) = \bar{4}+H = \bar{0}+H \leftarrow \text{identity element}$$

Thus,  $\bar{1}+H$  has order 4.

order of  $\bar{2}+H$

$$\bar{2}+H \neq \bar{0}+H$$

$$(\bar{2}+H) + (\bar{2}+H) = \bar{4}+H = \bar{0}+H \leftarrow \text{identity element}$$

Thus,  $\bar{2}+H$  has order 2.

---

(d)

Since  $\mathbb{Z}_{12}$  is abelian, from a problem in this HW set,  $\mathbb{Z}_{12}/H$  is abelian.

From part (c) we have that

$$\langle \bar{1}+H \rangle = \{ \bar{1}+H, \bar{2}+H, \bar{3}+H, \bar{4}+H \} = \mathbb{Z}_{12}/H.$$

Thus,  $\mathbb{Z}_{12}/H$  is cyclic with  $\bar{1}+H$  as a generator.

③

(a) From problem 1 we get that

$$D_8 = \{1, r, r^2, r^3, s, sr, sr^2, sr^3\}$$

$$H = \langle r^2 \rangle = \{1, r^2\}$$

left cosets :

$$H = \{1, r^2\} = r^2 H$$

$$rH = \{r, r^3\} = r^3 H$$

$$sH = \{s, sr^2\} = sr^2 H$$

$$srH = \{sr, sr^3\} = sr^3 H$$

So,

$$D_8/H = \{H, rH, sH, srH\}$$

The identity element is  $H = 1H$ .

---

(b)

$$(rH)(sH) = rsH = sr^{-1}H = sr^3H = srH$$

$$(srH)(srH) = (sr)(sr)H = ssr^{-1}r = s^2H = 1H = H$$



(c)

Since

$$(rH)(rH) = r^2H = H$$

We know that

$$(rH)^{-1} = rH$$

Since

$$(sH)(sH) = s^2H = 1H = H$$

We know that

$$(sH)^{-1} = H.$$

(d)

$$H \leftarrow \text{identity}$$

$H$  has order 1

$$sH \neq H$$

$$(sH)(sH) = s^2H = H \leftarrow \text{identity}$$

So,  $sH$  has order 2

$$rH \neq H$$

$$(rH)(rH) = r^2H = H$$

So,  $rH$  has order 2

identity  
↓

$$srH \neq H$$

$$(srH)(srH) = sr sr H = H$$

So,  $srH$  has order 2

see part (b)

identity

(e)

From (d) there are no elements of  $D_8/H$  that have order 4.

So no element will generate all of  $D_8/H = \{H, rH, sH, srH\}$

Thus,  $D_8/H$  is not cyclic.

---

(4)

(a)  $G = \mathbb{Z}_3 \times \mathbb{Z}_3 = \{(\bar{0}, \bar{0}), (\bar{0}, \bar{1}), (\bar{0}, \bar{2}), (\bar{1}, \bar{0}), (\bar{1}, \bar{1}), (\bar{1}, \bar{2}), (\bar{2}, \bar{0}), (\bar{2}, \bar{1}), (\bar{2}, \bar{2})\}$

$$H = \langle (\bar{0}, \bar{1}) \rangle = \{(\bar{0}, \bar{0}), (\bar{0}, \bar{1}), (\bar{0}, \bar{2})\}$$

left cosets:

$$\begin{aligned}(\bar{0}, \bar{0}) + H &= \{(\bar{0}, \bar{0}), (\bar{0}, \bar{1}), (\bar{0}, \bar{2})\} = (\bar{0}, \bar{1}) + H = (\bar{0}, \bar{2}) + H \\(\bar{1}, \bar{0}) + H &= \{(\bar{1}, \bar{0}), (\bar{1}, \bar{1}), (\bar{1}, \bar{2})\} = (\bar{1}, \bar{1}) + H = (\bar{1}, \bar{2}) + H \\(\bar{2}, \bar{0}) + H &= \{(\bar{2}, \bar{0}), (\bar{2}, \bar{1}), (\bar{2}, \bar{2})\} = (\bar{2}, \bar{1}) + H = (\bar{2}, \bar{2}) + H\end{aligned}$$

So,

$$G/H = \{(\bar{0}, \bar{0}) + H, (\bar{1}, \bar{0}) + H, (\bar{2}, \bar{0}) + H\}$$

The identity element is  $(\bar{0}, \bar{0}) + H$

---

(b)

$$\begin{aligned}[(\bar{1}, \bar{2}) + H] + [(\bar{1}, \bar{1}) + H] &= (\bar{2}, \bar{3}) + H = (\bar{2}, \bar{0}) + H \quad \begin{array}{c} \text{mod } 3 \\ \swarrow \quad \searrow \end{array} \\[(\bar{0}, \bar{1}) + H] + [(\bar{2}, \bar{1}) + H] &= (\bar{2}, \bar{2}) + H = (\bar{2}, \bar{0}) + H \quad \begin{array}{c} \uparrow \\ \text{from (a)} \end{array}\end{aligned}$$

(c)

Note that

$$(\bar{1}, \bar{2}) + H = (\bar{1}, \bar{0}) + H \leftarrow \text{from part (a)}$$

So we can instead find the inverse of  $(\bar{1}, \bar{0}) + H$ .

We have that:

$$[(\bar{1}, \bar{0}) + H] + [(\bar{2}, \bar{0}) + H] = (\bar{3}, \bar{0}) + H = (\bar{0}, \bar{0}) + H$$

mod 3

Thus, the inverse of  $(\bar{1}, \bar{2}) + H = (\bar{1}, \bar{0}) + H$  is  $(\bar{2}, \bar{0}) + H$ .

---

Now let's look at  $(\bar{2}, \bar{2}) + H$ .

From part (a) we get

$$(\bar{2}, \bar{2}) + H = (\bar{2}, \bar{0}) + H$$

And above we found that the inverse of  $(\bar{2}, \bar{0}) + H$  is  $(\bar{1}, \bar{0}) + H$ .

---

(d)

$(\bar{0}, \bar{0}) + H$  is the identity so it has order 1.

$(\bar{1}, \bar{0}) + H$   
 $[(\bar{1}, \bar{0}) + H] + [(\bar{1}, \bar{0}) + H] = (\bar{2}, \bar{0}) + H$   
 $[(\bar{1}, \bar{0}) + H] + [(\bar{1}, \bar{0}) + H] + [(\bar{1}, \bar{0}) + H] = (\bar{3}, \bar{0}) + H = (\bar{0}, \bar{0}) + H$   
 $(\bar{1}, \bar{0}) + H$  is the identity so it has order 3

} these are not the identity

$(\bar{2}, \bar{0}) + H$   
 $[(\bar{2}, \bar{0}) + H] + [(\bar{2}, \bar{0}) + H] = (\bar{4}, \bar{0}) + H = (\bar{1}, \bar{0}) + H$   
 $[(\bar{2}, \bar{0}) + H] + [(\bar{2}, \bar{0}) + H] + [(\bar{2}, \bar{0}) + H] = (\bar{6}, \bar{0}) + H = (\bar{0}, \bar{0}) + H$   
 $(\bar{2}, \bar{0}) + H$  is the identity so it has order 3

} not the identity

element	order
$(\bar{0}, \bar{0}) + H$	1
$(\bar{1}, \bar{0}) + H$	3
$(\bar{2}, \bar{0}) + H$	3

(e)  $G/H = \{(\bar{0}, \bar{0}) + H, (\bar{1}, \bar{0}) + H, (\bar{2}, \bar{0}) + H\}$

Thus,  $G/H$  has size 3.

The generators are the elements of order 3. There are two of them:

$$\left. \begin{array}{l} (\bar{1}, \bar{0}) + H \\ (\bar{2}, \bar{0}) + H \end{array} \right\} \begin{array}{l} \text{both of these} \\ \text{generate } G/H \end{array}$$

So,  $G/H$  is cyclic.

**(5)(a)** Let  $e$  be the identity element of  $G$ .

( $\Rightarrow$ ) Suppose that  $aH = bH$ .

Since  $H \leq G$  we know that  $e \in H$ .

Thus,  $a = ae \in aH$ .

recall that  
 $xH = \{xh \mid h \in H\}$

Since  $aH = bH$  and  $a \in aH$   
we know that  $a \in bH$ .

---

( $\Leftarrow$ ) Suppose that  $a \in bH$ .

Thus,  $a = bh$  where  $h \in H$ .

Let's show that  $aH = bH$ .

First we show that  $aH \subseteq bH$ .

Let  $c \in aH$ .

Then,  $c = ah_1$  where  $h_1 \in H$ .

Thus,  
 $c = ah_1 = bh_1 = b(\underbrace{hh_1}) \in bH$

Hence  $c \in bH$ .

Therefore,  $aH \subseteq bH$ .

this is in  $H$   
because  $H \leq G$   
and so  $H$  is closed  
under the group operation

Now we show that  $bH \subseteq aH$

Let  $d \in bH$ .

Then,  $d = bh_2$  where  $h_2 \in H$ .

So,

$$d = bh_2 = ah^{-1}h_2 = a(h^{-1}h_2) \in aH$$

↑  
 $a = bh$   
gives  
 $ah^{-1} = b$

$h^{-1} \in H$  since  $h \in H$  and  $H \leq G$   
 $h^{-1}h_2 \in H$  since  $h^{-1}, h_2 \in H$   
which gives  $h^{-1}h_2 \in H$   
since  $H \leq G$

Thus,  $d \in aH$

So,  $bH \subseteq aH$ .

Since  $aH \subseteq bH$  and  $bH \subseteq aH$  we  
know that  $aH = bH$ .



**(5)(b)** Just use part (a).

We have that

$$aH = bH \quad \left[ \text{from part (a)} \right]$$

$$\text{iff } a \in bH$$

$$\text{iff } a = bh \text{ for some } h \in H$$

since  
 $bH = \{bh \mid h \in H\}$





⑤(c)

Define  $f: H \rightarrow aH$  by  $f(h) = ah$ .

Let's show that  $f$  is one-to-one and onto.

$f$  is one-to-one:

Suppose  $f(h_1) = f(h_2)$ .

Then,  $ah_1 = ah_2$ .

So,  $a^{-1}ah_1 = a^{-1}ah_2$ .

Thus,  $h_1 = h_2$ .

Hence  $f$  is one-to-one.

$f$  is onto:

Let  $x \in aH$ .

Then,  $x = ah$  for some  $h \in H$ .

And  $f(h) = ah = x$

Thus,  $f$  is onto.

Since  $f$  is one-to-one and onto  
we know that  $H$  and  $aH$   
have the same size.

That is,  $|H| = |aH|$ .



⑥ Let  $|G| = pq$  where  $p$  and  $q$  are primes.  
Let  $H \leq G$ .

By Lagrange's theorem we know that  
 $|H|$  divides  $|G|$ .

Thus,  $|H| = 1, p, q$ , or  $pq$ .

since  $p$   
and  $q$  are  
primes

Since  $H \neq G$  we know  $|H| \neq pq$ .

Thus,  $|H| = 1, p$ , or  $q$ .

If  $|H| = 1$ , then  $H = \{e\}$ .

Then  $H = \langle e \rangle$  is cyclic.

If  $|H| = p$ , then by a theorem  
from class since  $p$  is prime  
we must have that  $H$  is cyclic.

If  $|H| = q$ , then by a theorem  
from class since  $q$  is prime  
we must have that  $H$  is cyclic.

In all three cases we have that  
 $H$  is cyclic.



⑦ Let  $|G| = n$ .

Suppose that  $x \in G$ .

Consider the subgroup

$$H = \langle x \rangle$$

generated by  $x$ .

Recall from class that the order of  $x$  is  $|\langle x \rangle| = |H|$ .

By Lagrange's theorem,  $|H|$  divides  $|G|$ .

Thus,  $n = |H| \cdot k$  for some integer  $k \geq 1$ .

Hence,

$$x^n = x^{|H| \cdot k} = (x^{|H|})^k = e^k = e$$

$|H|$  is the order of  $x$

Thus,  $x^n = e$



⑧(a) Suppose  $G$  is abelian.

Let  $x, y \in G/H$ .

Then,  $x = aH$  and  $y = bH$  for some  $a, b \in G$ .

So,

$$xy = (aH)(bH) = (ab)H = (ba)H = (bH)(aH) = yx$$

def of operation in  $G/H$

since  $G$  is abelian

Hence  $xy = yx$ .

So,  $G/H$  is abelian. ◻

⑧(b)

def of  $G/H$  operation

$$\text{Since } (aH)(\bar{a}'H) = a\bar{a}'H = eH = H$$

We know that  $(aH)^{-1} = \bar{a}'H$

identity element of  $G/H$

⑧ (<)

Let  $k \in \mathbb{Z}$ .

If  $k=0$ , then  $(aH)^0 = eH = a^0 H$

If  $k \geq 1$ , then

$$(aH)^k = \underbrace{(aH)(aH)\cdots(aH)}_{k \text{ times}} = a^k H$$

def of  $G/H$  operation

If  $k \leq -1$ , then

$$(aH)^k = \underbrace{(aH)^{-1}(aH)^{-1}\cdots(aH)^{-1}}_{-k \text{ times}}$$

part (b)

$$= \underbrace{(\bar{a}^{-1}H)(\bar{a}^{-1}H)\cdots(\bar{a}^{-1}H)}_{-k \text{ times}}$$

$$= (\bar{a}^{-1})^{-k} H$$

$$= a^k H.$$

We have show that  $(aH)^k = a^k H$  for all the cases of  $k$ .



⑧(d) Let  $G$  be cyclic.

Then,  $G = \langle g \rangle$  for some  $g \in G$ .

Let's show that  $gH$  generates  $G/H$ .

Let  $x \in G/H$ .

Then  $x = aH$  for some  $a \in G$ .

Since  $a \in G$  and  $G = \langle g \rangle$  we have that  $a = g^k$  for some  $k \in \mathbb{Z}$ .

Therefore,

$$x = aH = g^k H = \underset{\substack{\uparrow \\ \text{part (c)}}}{gH}^k$$

Hence,  $G/H = \langle gH \rangle$ .

So,  $G/H$  is cyclic.



⑨ Suppose that  $H \trianglelefteq G$  and  $K \trianglelefteq G$ .

We showed that  $H \cap K \leq G$  is HW 2.

So we just have to show that  $H \cap K$  is normal.

Let  $g \in G$  and  $a \in H \cap K$ .

We need to show that  $gag^{-1} \in H \cap K$ .

Since  $a \in H \cap K$  we know that  $a \in H$  and  $a \in K$ .

Since  $g \in G$  and  $a \in H$  and  $H$  is normal we know that  $gag^{-1} \in H$ .

Since  $g \in G$  and  $a \in K$  and  $K$  is normal we know that  $gag^{-1} \in K$ .

Thus,  $gag^{-1} \in H \cap K$ .

Therefore,  $H \cap K \trianglelefteq G$ .

From class:  
Let  $N \leq G$ .  
Then  $N$  is normal  
iff  
 $gng^{-1} \in N$   
for all  
 $n \in N$   
and  
 $g \in G$



⑩  $\varphi: G_1 \rightarrow G_2$  is a homomorphism.

Let  $e_1$  and  $e_2$  be the identity elements of  $G_1$  and  $G_2$  respectively.

Recall that

$$\ker(\varphi) = \{x \mid x \in G_1 \text{ and } \varphi(x) = e_2\}$$

From HW 4 we know that

$\ker(\varphi)$  is a subgroup of  $G_1$ .

Let's show that  $\ker(\varphi)$  is normal.

Let  $H = \ker(\varphi)$ .

Let  $g \in G$  and  $h \in H$ .

We need to show that  $ghg^{-1} \in H$ .

Since  $h \in H$  and  $H = \ker(\varphi)$   
we know that  $\varphi(h) = e_2$

Thus,

$$\begin{aligned}\varphi(ghg^{-1}) &= \varphi(g)\varphi(h)\varphi(g^{-1}) \\ &= \varphi(g)e_2\varphi(g^{-1}) \\ &= \varphi(g)\varphi(g^{-1}) \\ &= \varphi(gg^{-1})\end{aligned}$$

since  $\varphi$  is a homomorphism

From class  
Let  $H \leq G$ .

Then:  
 $H$  is normal  
iff  
 $ghg^{-1} \in H$   
for all  
 $h \in H$   
and  
 $g \in G$



$$= \varphi(e_1)$$


Hw 4

$$= e_2$$

Thus,  $\varphi(ghg^{-1}) = e_2$ .

So,  $ghg^{-1} \in \ker(\varphi)$ .

Hence  $\ker(\varphi) \trianglelefteq G_1$ .



⑪ Let  $H \leq G$  and  $G$  be finite.  
From class, to show that  $H \trianglelefteq G$  we  
can show that  $gHg^{-1} = H$  for  
all  $g \in G$  where  
$$gHg^{-1} = \{ ghg^{-1} \mid h \in H \}$$

Let  $g \in G$ .

Let's show that  $gHg^{-1} = H$ .

Define  $\varphi_g: H \rightarrow G$  by  $\varphi_g(h) = ghg^{-1}$ .

Note that  $\varphi_g$  is a homomorphism because

$$\begin{aligned}\varphi_g(h_1 h_2) &= gh_1 h_2 g^{-1} \\ &= gh_1 g^{-1} g h_2 g^{-1} \\ &= \varphi_g(h_1) \varphi_g(h_2).\end{aligned}$$

Note that

$$\begin{aligned}\text{im}(\varphi_g) &= \{ \varphi_g(h) \mid h \in H \} \\ &= \{ ghg^{-1} \mid h \in H \}\end{aligned}$$

$$= gHg^{-1}$$

Also note that  $\varphi_g$  is one-to-one because if  $\varphi_g(a) = \varphi_g(b)$  then  $ga\bar{g}^{-1} = gb\bar{g}^{-1}$  and so  $g^{-1}(ga\bar{g}^{-1})g = g^{-1}(gb\bar{g}^{-1})g$  which gives  $a=b$ .

Also recall from Hw 4 / class that  $\text{im}(\varphi_g) \leq G$ .

Thus,  $gHg^{-1} \leq G$ .

Since  $\varphi_g$  is one-to-one we get that  $|H| = |\text{im}(\varphi_g)| = |gHg^{-1}|$ .

Since  $H$  is the only subgroup of  $G$  of size  $|H|$  and  $gHg^{-1} \leq G$  we know that  $H = gHg^{-1}$ .

Thus,  $H$  is normal.

