

Math 4300 - Homework # 4

Betweenness

-
1. In the Euclidean plane, let $A = (-1, -2)$, $B = (2, 1)$, and $C = (0, -1)$.
 - (a) Determine if A, B, C are collinear or not. Draw a picture.
 - (b) If the points are collinear, Determine if $A - B - C$, $A - C - B$, or $B - A - C$.
 - (c) Determine if $B - C - A$.
-
2. In the hyperbolic plane, let $A = (1, 2)$, $B = (3, 4)$ and $C = (4, \sqrt{19})$.
 - (a) Determine if A, B, C are collinear or not. Draw a picture.
 - (b) If the points are collinear, Determine if $A - B - C$, $A - C - B$, or $B - A - C$.
 - (c) Determine if $B - C - A$.
-
3. In the hyperbolic plane, let $A = (1, 2)$, $B = (1, 4)$ and $C = (1, 5)$.
 - (a) Determine if A, B, C are collinear or not. Draw a picture.
 - (b) If the points are collinear, Determine if $A - B - C$, $A - C - B$, or $B - A - C$.
-
4. Let $(\mathcal{P}, \mathcal{L}, d)$ be a metric geometry. Let $A, B \in \mathcal{P}$ with $A \neq B$.
Let $C \in \overleftrightarrow{AB}$. Prove that one and only one of the following can be true:
 $C - A - B$ or $C = A$ or $A - C - B$ or $C = B$ or $A - B - C$.
-
5. Let $(\mathcal{P}, \mathcal{L}, d)$ be a metric geometry. Let ℓ be a line and A, B, C be distinct points on ℓ . Prove that either $A - B - C$ or $A - C - B$ or $B - A - C$.
-

6. Let $(\mathcal{P}, \mathcal{L}, d)$ be a metric geometry. Let A, B, C, D be points from \mathcal{P} .
Prove that if $A - B - C$ and $B - C - D$, then $A - B - D$ and $A - C - D$.
-

7. Let $(\mathcal{P}, \mathcal{L}, d)$ be a metric geometry. Let A, B, C, D be points from \mathcal{P} .
Prove that if $A - C - D$ and $A - C - B$, then $A - D - B$ or $A - B - D$.
-

8. Let $(\mathcal{P}, \mathcal{L}, d)$ be a metric geometry. Let A, B, C, D be points from \mathcal{P} .
Prove that if $A - D - C$ and $A - C - B$, then $A - D - B$.
-

9. Let $(\mathcal{P}, \mathcal{L}, d)$ be a metric geometry. Let A, B, P, Q be points from \mathcal{P} .
Prove that if $A - Q - B$ and $A - P - B$ and $P - C - Q$, then $A - C - B$.
-

10. Consider the Euclidean plane $\mathcal{E} = (\mathbb{R}^2, \mathcal{L}_E, d_E)$. Let $A, B, C \in \mathbb{R}^2$ be distinct points. Prove that $A - B - C$ if and only if there exists a real number t with $0 < t < 1$ and $B = A + t(C - A)$.
-