

# Math 5680

## Homework # 2

### Sequences of functions, series of functions, Weierstrass M-Test, Analytic convergence theorem

---

1. Let  $A = \mathbb{R} \subseteq \mathbb{C}$  and  $f_n : A \rightarrow \mathbb{C}$  be defined for  $n \geq 2$  as follows

$$f_n(x) = \begin{cases} -1 & \text{for } x \leq -1/n \\ nx & \text{for } -1/n < x < 1/n \\ 1 & \text{for } 1/n \leq x \end{cases}$$

- (a) Draw a picture of  $f_n$  for  $n = 2$  and  $n = 3$  and  $n = 4$ .  
(b) Let

$$f(x) = \begin{cases} -1 & \text{for } x < 0 \\ 0 & \text{for } x = 0 \\ 1 & \text{for } 0 < x \end{cases}$$

Prove that  $f_n$  converges pointwise to  $f$  on  $A = \mathbb{R}$ .

---

2. Let

$$g(z) = \sum_{n=1}^{\infty} \frac{z^n}{n}$$

Given a positive real number  $r$ , define the sets

$$A_r = \{z \mid |z| \leq r\}$$

- (a) Draw a picture of  $A_1$  and  $A_\pi$ .  
(b) If  $0 \leq r < 1$  prove that that the series  $g(z)$  converges absolutely and uniformly on  $A_r$ .
- 

3. Let

$$g(z) = \sum_{n=1}^{\infty} \frac{1}{z^n}$$

and  $A = \{z \mid |z| > 1\}$ .

- (a) Show that  $g(z)$  is analytic on  $A$ .  
(b) Find the derivative of  $g(z)$  on  $A$ .
- 

4. Let

$$g(z) = \sum_{n=1}^{\infty} \frac{1}{n!} z^n$$

- (a) Show that  $g$  is analytic on  $A = \mathbb{C} - \{0\}$ .  
(b) Find the derivative of  $g(z)$  on  $A$ .
- 

5. Suppose that  $\sum_{k=1}^{\infty} g_k(z)$  converges uniformly on some subset  $A$  of  $\mathbb{C}$ . Prove that the sequence  $(g_k)$  converges uniformly to the zero function  $f_0$  on  $A$ . Here  $f_0(z) = 0$  for all  $z \in A$ .
- 

### THE NEXT PROBLEM ISN'T NECESSARY TO DO.

We used the fact given below in the proof of the analytic convergence theorem. Do the problem if you want an extra challenge and you are familiar with theorems about compactness.

A. Let  $A \subseteq \mathbb{C}$  be an open set and  $z_0 \in A$ . Let  $r > 0$ . Suppose that

$$B = \{z \mid |z - z_0| \leq r\}$$

is contained in  $A$ . Prove that there is a number  $\rho > r$  such that the circle  $\gamma$  of radius  $\rho$  centered at  $z_0$  satisfies (i)  $\gamma$  is contained in  $A$  and (ii)  $\gamma$  contains  $B$ .

[**Hint:** The boundary of  $B$  is a compact subset of  $\mathbb{C}$ . For each point on the boundary of  $B$  pick a open disk that surrounds that point and is contained in  $A$ . Then shrink those disks in half. Then use compactness to get a finite sub-cover.]