

4 **UPPER BOUNDS FOR THE STRONG CHROMATIC INDEX**  
5 **OF HALIN GRAPHS**

6 ZIYU HU

7 *Department of Mathematical Sciences, Florida Atlantic University*

8 **e-mail:** azuth.hu@gmail.com

9 KO-WEI LIH

10 *Institute of Mathematics, Academia Sinica*

11 **e-mail:** makwlih@sinica.edu.tw

12 AND

13 DAPHNE DER-FEN LIU <sup>1</sup>

14 *Department of Mathematics, California State University Los Angeles*

15 **e-mail:** dliu@calstatela.edu

16 **Abstract**

17 The strong chromatic index of a graph  $G$ , denoted by  $\chi'_s(G)$ , is the  
18 minimum number of vertex induced matchings needed to partition the edge  
19 set of  $G$ . Let  $T$  be a tree without vertices of degree 2 and have at least  
20 one vertex of degree greater than 2. We construct a Halin graph  $G$  by  
21 drawing  $T$  on the plane and then drawing a cycle  $C$  connecting all its leaves  
22 in such a way that  $C$  forms the boundary of the unbounded face. We call  
23  $T$  the characteristic tree of  $G$ . Let  $G$  denote a Halin graph with maximum  
24 degree  $\Delta$  and characteristic tree  $T$ . We prove that  $\chi'_s(G) \leq 2\Delta + 1$  when  
25  $\Delta \geq 4$ . In addition, we show that if  $\Delta = 4$  and  $G$  is not a wheel, then  
26  $\chi'_s(G) \leq \chi'_s(T) + 2$ . A similar result for  $\Delta = 3$  was established by Lih and  
27 Liu [25].

28 **Keywords:** Strong edge-coloring, strong chromatic index, Halin graphs.

29 **2010 Mathematics Subject Classification:** 05C15.

---

<sup>1</sup>Supported in part by grants NASA MIRO NX15AQ06A and NSF DMS 1600778.

30

## 1. INTRODUCTION

31 Let  $G$  be a simple graph. The *distance* between two edges  $e$  and  $e'$  in  $G$  is the  
 32 minimum  $k$  for which there is a sequence  $e = e_0, e_1, \dots, e_k = e'$  of distinct edges  
 33 such that for  $1 \leq i \leq k$ ,  $e_{i-1}$  and  $e_i$  share an end vertex. A *strong edge-coloring*  
 34 of a graph is a function that assigns to each edge a color such that any two edges  
 35 with distance at most two must receive different colors. A *strong  $k$ -edge-coloring*  
 36 is a strong edge-coloring using  $k$  colors. The *strong chromatic index* of a graph  
 37  $G$ , denoted by  $\chi'_s(G)$ , is the minimum  $k$  such that  $G$  admits a strong  $k$ -edge-  
 38 coloring. The pre-image of each color in a strong edge-coloring is an induced  
 39 matching. Thus, the strong chromatic index is also the minimum number of  
 40 vertex induced matchings needed to partition the edge set of  $G$ .

41 Denote the maximum degree of a graph  $G$  by  $\Delta(G)$  (or, simply by  $\Delta$  when  
 42  $G$  is clear in the context). A trivial upper bound is that  $\chi'_s(G) \leq 2\Delta(G)^2 -$   
 43  $2\Delta(G) + 1$ . Fouquet and Jolivet [16] established a Brooks type upper bound  
 44  $\chi'_s(G) \leq 2\Delta(G)^2 - 2\Delta(G)$ , which is not true only for  $G = C_5$  as pointed out by  
 45 Shiu and Tam [31]. The following conjecture was posed by Erdős and Nešetřil  
 46 [13, 14]:

47 **Conjecture 1.** *For any graph  $G$  of maximum degree  $\Delta$ ,*

$$48 \quad \chi'_s(G) \leq \begin{cases} \frac{5}{4}\Delta^2 & \text{if } \Delta \text{ is even;} \\ \frac{5}{4}\Delta^2 - \frac{1}{2}\Delta + \frac{1}{4} & \text{if } \Delta \text{ is odd.} \end{cases}$$

49 For graphs with maximum degree  $\Delta(G) = 3$ , Conjecture 1 was verified by An-  
 50 dersen [1] and by Horák, Qing and Trotter [22], independently. For  $\Delta(G) = 4$ ,  
 51 while Conjecture 1 asserts that  $\chi'_s(G) \leq 20$ , Horák [21] obtained  $\chi'_s(G) \leq 23$  and  
 52 Cranston [11] proved  $\chi'_s(G) \leq 22$ . For general graphs  $G$  with maximum degree  $\Delta$ ,  
 53 Molloy and Reed [28] showed that  $\chi'_s(G) \leq 1.998\Delta^2$ . Most recently, this bound  
 54 has been improved by Bruhn and Joos [6] to  $1.93\Delta^2$ .

55 Strong edge-coloring for planar graphs has been investigated by many au-  
 56 thors. Fouquet and Jolivet [16, 17] first studied strong edge-coloring for cubic  
 57 planar graphs. Let  $G$  be a planar graph with maximum degree  $\Delta$  and girth  $g$ .  
 58 Faudree et al. [15] proved that  $\chi'_s(G) \leq 4\Delta + 4$ . Bensmail et al. [2] established  
 59 the bound  $\chi'_s(G) \leq 3\Delta + 1$  for  $g \geq 6$ . Hudák et al. [23] showed  $\chi'_s(G) \leq 3\Delta$  if  
 60  $g \geq 7$ , and the bound is sharp for some subcubic (that is,  $\Delta \leq 3$ ) planar graphs.  
 61 Furthermore, Hocquard et al. [19] showed that  $\chi'_s(G) \leq 9$  for subcubic planar  
 62 graphs  $G$  which do not contain cycles of lengths 4 or 5. DeOrsey et al. [12] re-  
 63 cently reduced this bound to  $\chi'_s(G) \leq 5$  if  $g \geq 30$ . For planar graphs with large  
 64 girth, Borodin and Ivanova [3] established a rather tight bound  $\chi'_s(G) \leq 2\Delta - 1$   
 65 if  $g \geq 40\lfloor \Delta/2 \rfloor + 1$ ; Chang et al. [10] further confirmed that the bound also holds  
 66 if  $g \geq 10\Delta + 46$ . Clearly, the bound  $\chi'_s(G) \leq 2\Delta - 1$  becomes sharp when  $G$   
 67 contains two adjacent vertices of maximum degree  $\Delta$ .

68 By definition, a trivial lower bound of  $\chi'_s(G)$  for a graph  $G$  would be  $\sigma(G)$ ,  
69 where

$$70 \quad \sigma(G) := \max\{\deg(u) + \deg(v) - 1 \mid uv \in E(G)\}.$$

71 If  $G$  has no edges, then define  $\sigma(G) = 0$ . It is known and easy to verify that  
72 for a tree  $T$ , we have  $\chi'_s(T) = \sigma(T)$ . Wu and Lin [32] proved that if  $\sigma(G) \leq 4$   
73 and  $G$  is not isomorphic to the graph of the 5-cycle with a chord connecting two  
74 non-adjacent vertices, then  $\chi'_s(G) \leq 6$ . Recently, Chang and Duh [8] assert that  
75  $\chi'_s(G) = \sigma(G)$  if  $G$  is a planar graph with  $\sigma(G) = \sigma \geq 5$ ,  $\sigma \geq \Delta(G) + 2$ , and girth  
76  $g \geq 5\sigma + 16$ . This result implies that a planar graph with large girth behaves like  
77 a tree locally.

78 A Halin graph is a plane graph  $G$  constructed as follows. Let  $T$  be a tree with  
79 at least 4 vertices, called the *characteristic tree* of  $G$ . All vertices of  $T$  are either  
80 of degree 1, called leaves, or of degree at least 3. We draw  $T$  on the plane. Let  $C$   
81 be a cycle, called the *adjoint cycle* of  $G$ , connecting all leaves of  $T$  in such a way  
82 that  $C$  forms the boundary of the unbounded face. We usually write  $G = T \cup C$   
83 to reveal the characteristic tree and the adjoint cycle. For  $n \geq 3$ , the wheel  $W_n$   
84 with  $n + 1$  vertices is a particular Halin graph whose characteristic tree is the  
85 complete bipartite graph  $K_{1,n}$  (called a *star*). A graph is said to be cubic if the  
86 degree of every vertex is 3. For  $h \geq 1$ , a cubic Halin graph  $Ne_h$ , called a *necklace*,  
87 was introduced in [30]. Its characteristic tree  $T$  consists of the path  $v_0, v_1, \dots,$   
88  $v_h, v_{h+1}$  and leaves  $v'_1, v'_2, \dots, v'_h$  such that the unique neighbor of  $v'_i$  in  $T$  is  $v_i$   
89 for  $1 \leq i \leq h$  and vertices  $v_0, v'_1, \dots, v'_h, v_{h+1}$  are connected in this order to form  
90 the adjoint cycle  $C_{h+2}$ .

91 Lai, Lih and Tsai [24] proved the following result:

92 **Theorem 2** [24]. *If a Halin graph  $G = T \cup C$  is different from a certain necklace*  
93  *$Ne_2$  and any wheel  $W_n$ ,  $n \not\equiv 0 \pmod{3}$ , then  $\chi'_s(G) \leq \chi'_s(T) + 3$ .*

94 For cubic Halin graphs, Lih and Liu improved the above bound as follows:

95 **Theorem 3** [25]. *A cubic Halin graph  $G$  different from  $Ne_2$  or  $Ne_4$  satisfies*  
96  *$\chi'_s(G) \leq 7$ .*

97 The exact values of  $\chi'_s(G)$  for special families of cubic Halin graphs were deter-  
98 mined by Shiu and Tam [31] and by Chang and Liu [9].

99 For a Halin graph  $G = T \cup C$  with maximum degree  $\Delta$ , since  $\chi'_s(T) \leq 2\Delta - 1$ ,  
100 the bound in Theorem 2 implies that  $\chi'_s(G) \leq 2\Delta + 2$ . We improve this bound  
101 and establish a similar result of Theorem 3 for Halin graphs of maximum degree  
102 4.

103 **Theorem 4.** *Let  $G$  be a Halin graph with maximum degree  $\Delta \geq 4$ . Then  $\chi'_s(G) \leq$*   
104  *$2\Delta + 1$ .*

105 **Theorem 5.** *Let  $G = T \cup C$  be a Halin graph with maximum degree  $\Delta = 4$ , and*  
 106 *let  $G$  be different from a wheel. Then  $\chi'_s(G) \leq \chi'_s(T) + 2$ .*

107 Both bounds in Theorems 4 and 5 are sharp. Consider the graph  $G$  in  
 108 Figure 1. A strong edge-coloring of  $G$  must use at least 7 colors on the edges  
 109 incident to  $u$  or  $v$ . Let these colors be  $\{1, 2, \dots, 7\}$ . Next, since the edges  $w_1$   
 110 and  $w_2$  must use colors different from  $\{1, 2, \dots, 7\}$ , at least 8 colors are needed.  
 111 Assume we only have 8 colors. Then  $w_1$  and  $w_2$  must be colored by the same new  
 112 color, say color 8. This implies that the four edges  $e_1, e_2, e_3, e_4$  shown in Figure 1  
 113 only have three admissible colors, from the set  $\{5, 6, 7\}$ , which is a contradiction  
 114 as these edges must receive different colors. Hence  $\chi'_s(G) \geq 9$ . By coloring  $e_1, e_2,$   
 115  $e_3, e_4$  with colors 5, 6, 7, 9 and the last edge with color 4 it follows that  $\chi'_s(G) = 9$ .  
 116 This example shows that both bounds in Theorems 4 and 5 are sharp.




Figure 1.: An example showing sharp bounds of Theorems 4 and 5.

117

## 2. PROOF OF THEOREM 4

118 A *double star* is a tree with exactly two non-leaf vertices. Denote  $D_{a,b}$  a double  
 119 star where  $a \leq b$  are the degrees of the two non-leaf vertices. Prior to the proof  
 120 of Theorem 4, we quote several known results as follows.

121 **Lemma 6** [24]. *Let  $G = T \cup C$  be a Halin graph. If  $T = D_{a,b}$  is a double star*  
 122 *with  $a \leq b$ , then*

$$123 \quad \chi'_s(G) = \begin{cases} \chi'_s(T) + 4 & \text{if } a = b = 3; \\ \chi'_s(T) + 2 & \text{if } a = 3 \text{ and } b \geq 4; \\ \chi'_s(T) + 1 & \text{if } a \geq 4. \end{cases}$$

124 *If  $T = K_{1,k}$  (that is,  $G$  is a wheel  $W_k$ ), then*

$$125 \quad \chi'_s(W_k) = \begin{cases} k + 3 & \text{if } k \equiv 0 \pmod{3}; \\ k + 5 & \text{if } k = 5; \\ k + 4 & \text{otherwise.} \end{cases}$$

126 **Lemma 7** [30]. *Suppose  $h \geq 1$ . Then*

$$127 \quad \chi'_s(Ne_h) = \begin{cases} 6 & \text{if } h \text{ is odd;} \\ 7 & \text{if } h \geq 6 \text{ and } h \text{ is even;} \\ 8 & \text{if } h = 4; \\ 9 & \text{if } h = 2. \end{cases}$$

128 **Proof of Theorem 4.** Let  $G = T \cup C$  be a Halin graph with  $\Delta(G) \geq 4$ . If  $T$  is a  
 129 star or a double star, by Lemma 6, the conclusion of Theorem 4 follows. Assume  
 130 that  $T$  is neither a star nor a double star. We proceed by induction on  $|C|$ , the  
 131 length of  $C$ . The shortest length of  $C$  is 6. Three possible graphs along with  
 132 their corresponding strong edge-colorings satisfying the desired upper bounds are  
 133 shown in Figure 2. So the result follows.




Figure 2.: All Halin graphs with  $|C| = 6$  and  $\Delta(G) = 4$ .

134 Assume  $|C| \geq 7$ . Let  $P = u_0, u_1, \dots, u_l$  be a longest path in  $T$  with length  $l$ .  
 135 As  $T$  is neither a star nor a double star, so  $l \geq 4$ . Without loss of generality, we  
 136 assume  $\deg_G(u_{l-1}) \geq \deg_G(u_1)$ .

137 Denote  $u_1 = v$ ,  $u_2 = u$ ,  $u_3 = w$ , and label the  $k \geq 2$  leaf neighbors of  $v$  as  
 138  $v_1, v_2, \dots, v_k$ . Since  $P$  is a longest path in  $T$ , it is easy to see that  $v_1, v_2, \dots, v_k$   
 139 must be on the adjoint cycle  $C$ . Let  $x_1, x_2, y_1, y_2$  be vertices on  $C$ , where  $x_1$  is  
 140 adjacent to  $v_1$  and  $x_2$ ;  $y_1$  is adjacent to  $v_k$  and  $y_2$ . Let  $x_3$  and  $y_3$  be vertices not  
 141 on  $C$ , where  $x_1x_3$  and  $y_1y_3$  are edges in  $T$  (see Figure 3).

142 Since  $G$  is a Halin graph and  $u$  is a vertex of degree at least 3, there exists a  
 143 path  $P'$  in  $T$  from  $u$  to  $x_1$  or from  $u$  to  $y_1$  with  $P \cap P' = \{u\}$ . Without loss of  
 144 generality, we shall assume that  $P'$  is from  $u$  to  $y_1$ . By our assumption that  $P$  is  
 145 a longest path, it must be that  $|P'| \leq 2$ . Thus, either  $u = y_3$  or  $u$  is adjacent to  
 146  $y_3$ .

147 In the following, we denote by  $G' = T' \cup C'$  the Halin graph obtained by  
 148 adding some new edges to an induced subgraph of  $G$  such that  $|C'| < |C|$  and  
 149  $\Delta(G') \leq \Delta(G)$ . If  $\Delta(G') \geq 4$  then  $\chi'_s(G') \leq 2\Delta(G) + 1$  holds because  $T'$  is a star  
 150 or double star (see the beginning of the proof) or by the inductive hypothesis as  
 151  $|C'| < |C|$ . If  $\Delta(G') = 3$  then  $\chi'_s(G') \leq 9 \leq 2\Delta(G) + 1$  by Theorem 2, Lemma




Figure 3.: The neighborhood around one end of the longest path  $P$ .

152 6, and because  $\Delta(G) \geq 4$ . In the following case analysis these steps will be  
 153 repeatedly used, while may not be mentioned explicitly all the time.

154 We call  $G'$  a *reduction* of  $G$ . Depending on various situations, different types  
 155 of  $G'$  are created. In the corresponding figures, the dashed lines represent new  
 156 edges added in  $G'$ , and dark vertices represent the vertices that are temporarily  
 157 deleted from  $G$ .

158 Let  $\psi$  be a strong edge-coloring of  $G'$  using the minimum number of colors.  
 159 A strong edge-coloring  $\phi$  of  $G$  is obtained as follows. We color the edges that are  
 160 in both  $G$  and  $G'$  by the same colors used in  $\psi$ , i.e., let  $\phi(e) = \psi(e)$  for every  
 161  $e \in E(G) \cap E(G')$ . For edges in  $e \in E(G) \setminus E(G')$ , we develop different coloring  
 162 schemes for different cases, and in each case, we give a strong edge-coloring  $\phi$  for  
 163  $G$  with at most  $2\Delta(G) + 1$  colors.

164 Case A.  $\deg_G(v) = 3$  There are three possibilities to consider.

165 A.1.  $u = y_3$ . Obtain the reduction  $G'$  of  $G$  by adding two new edges  $vx_1$  and  $vy_1$   
 166 to the induced subgraph of  $G$  on the vertex set  $V(G) \setminus \{v_1, v_2\}$ , as indicated in  
 167 Figure 4. Clearly,  $\Delta(G') = \Delta(G) \geq 4$  and  $|C'| < |C|$ .




Figure 4.: Case A.1.

168 Without loss of generality, assume that  $\psi(vx_1) = 1$  and  $\psi(vy_1) = 2$ . Let  
 169  $\phi(v_1x_1) = 1$  and  $\phi(v_2y_1) = 2$ . See Figure 4. We find admissible colors  $w_1$ ,  
 170  $w_2$ , and  $w_3$ , one by one. The colors that can not be assigned to  $vv_1$  are from  
 171  $\{1, 2, t_1, t_2\}$  and the labels used by edges incident to  $u$ . Therefore, there are  
 172 at most  $\Delta(G) + 4$  forbidden colors for  $vv_1$ . Since  $\Delta(G) \geq 4$ , there exists an  
 173 admissible color for  $vv_1$ . Color  $vv_1$  by such an admissible color  $w_1$ .

174 Next we color  $vv_2$  which has the forbidden colors in  $\{1, 2, w_1, s\}$  and the  
 175 labels used for edges incident to  $u$ . Similarly, we can find an admissible color for  
 176  $vv_2$ . Finally, the forbidden colors for  $v_1v_2$  are in  $\{1, 2, w_1, w_2, r_1, r_2, s, t_1, t_2\}$ . If  
 177  $s \in \{t_1, t_2\}$ , then there is an admissible color for  $v_1v_2$ . Otherwise, we re-color  $vv_1$   
 178 by  $s$ , creating an admissible color for  $v_1v_2$ .

179 A.2.  $u$  is adjacent to  $y_3$ , and  $\Delta(G) \geq 5$ . Obtain the reduction  $G'$  in the same way  
 180 as Case A.1, as indicated in Figure 5. Clearly,  $\Delta(G') = \Delta(G) \geq 4$  and  $|C'| < |C|$ .




Figure 5.: Case A.2.

181 Without loss of generality, assume that  $\psi(vx_1) = 1$  and  $\psi(vy_1) = 2$ . Let  
 182  $\phi(v_1x_1) = 1$  and  $\phi(v_2y_1) = 2$  (see Figure 5). We find admissible colors  $w_1$ ,  $w_2$ ,  
 183 and  $w_3$ , one by one. By the same argument in Case A.1, one can easily show that  
 184 there exists an admissible color  $w_1$ . Color  $vv_1$  by such an admissible color.

185 Next we color  $vv_2$  which has the forbidden colors in  $\{1, 2, w_1, s_1, s_2\}$  and the  
 186 labels used for edges incident to  $u$ . Since  $\Delta(G) \geq 5$ , we can find an admissible  
 187 color  $w_2$ . Finally, the forbidden colors for  $v_1v_2$  are in  $\{1, 2, w_1, w_2, r, s_1, s_2, t_1, t_2\}$ .  
 188 Thus, there exists an admissible color  $w_3$ .

189 A.3.  $u$  is adjacent to  $y_3$ , and  $\Delta(G) = 4$ . Then  $\deg_G(y_3)$  is either 3 or 4. Obtain  
 190 the reduction  $G'$  from  $G$  with partial labels to some vertices as indicated in Fig-  
 191 ure 6(a) and 6(b), respectively. Clearly,  $\Delta(G') \leq \Delta(G)$  and  $|C'| < |C|$ . Assume  
 192 that  $\deg_G(y_3) = 3$ . Then  $\Delta(G') = \Delta(G) = 4$ . We find admissible colors  $w_1$ ,  $w_2$ ,  
 193 and  $w_3$ , one after another. For  $v_1v_2$ , the forbidden colors are in  $\{1, 2, 3, r_1, t_1, t_2\}$ .  
 194 Hence there is an admissible color  $w_1$  for  $v_1v_2$ . Next, the forbidden colors for  
 195  $y_1y_2$  are in  $\{1, 2, 3, w_1, r_2, s_1, s_2\}$ . We can color  $y_1y_2$  by an admissible color  $w_2$ .

196 Finally, the forbidden colors for  $v_2y_1$  are in  $\{1, 2, 3, w_1, w_2, r_1, r_2\}$ . Again, there  
 197 exists an admissible color  $w_3$  for  $v_2y_1$ .




Figure 6.: Case A.3.

198 Assume  $\deg_G(y_3) = 4$ . Note, even if  $\Delta(G') = 3$  or  $T'$  is a star (or double  
 199 star), we can still find a strong edge coloring for  $G'$  by up to 9 colors. The  
 200 forbidden colors for  $y_1y_3$  are in  $\{1, 2, 3\}$  and labels used on edges incident to  $u$ .  
 201 Thus there are at most  $\Delta(G)+3$  forbidden colors. We color  $y_1y_3$  by an admissible  
 202 color  $w_1$ . Next, the forbidden colors for  $v_1v_2$  are  $\{1, 2, 3, 4, w_1, t_1, t_2\}$ . Because  
 203  $2\Delta(G) + 1 \geq 9$ , we can find an admissible color  $w_2$  for  $v_1v_2$ . The forbidden colors  
 204 for  $y_2z$  are in  $\{1, 2, 3, 4, w_1, r, s_1, s_2\}$ . Again, there is an admissible color  
 205  $w_3$  for  $y_2z$ . Finally, the forbidden colors for  $v_2y_1$  are from  $\{1, 2, 3, 4, w_1, w_2, w_3, r\}$ . So  
 206 there is an admissible color  $w_4$  for  $v_2y_1$ .

207 Case B.  $\deg_G(v) \geq 4$  We consider two cases separately.

208 B.1.  $\Delta(G) = 4$ . Then  $\deg_G(v) = 4$ . There are two subcases.

209 Subcase B.1.1.  $\deg_G(u) = 3$ . Obtain the reduction  $G'$  of  $G$  by adding two new  
 210 edges  $vx_1$  and  $vy_1$  to the induced subgraph of  $G$  on the vertex set  $V(G) \setminus$   
 211  $\{v_1, v_2, v_3\}$  as depicted in Figure 7.

212 Since we assumed earlier that  $\deg_G(u_{l-1}) \geq \deg_G(u_1) = \deg_G(v) = 4$ , we  
 213 have  $\Delta(G') = \Delta(G) = 4$ , and  $|C'| < |C|$  holds. We fix colors on some edges as  
 214 shown on Figure 7. Note that in Figure 7(a) we assign  $\phi(y_1y_2) = \phi(vv_2) = 3$   
 215 but in Figure 7(b) we assign  $\phi(y_1y_3) = \phi(vv_2) = 3$  and  $\phi(y_1y_2) = s$ . We find  
 216 admissible colors  $w_1, w_2, w_3$ , and  $w_4$ .

217 For the subcase depicted in Figure 7(a), the forbidden colors for  $vv_1$  are in  
 218  $\{1, 2, 3, t_1, t_2\}$  and the three colors used in the neighborhood of  $u$ . Thus, there  
 219 are at most 8 forbidden colors, implying there is an admissible color  $w_1$  for  $vv_1$ .  
 220 Next, the forbidden colors for  $vv_3$  are in  $\{1, 2, 3, w_1\}$  and the three colors used in  
 221 the neighborhood of  $u$ . There is an admissible color  $w_2$  for  $vv_3$ . The forbidden  
 222 colors for  $v_1v_2$  are in  $\{1, 2, 3, w_1, w_2, r_1, t_1, t_2\}$ , so there is an admissible color  $w_3$






Figure 7.: Subcase B.1.1.

223 for  $v_1v_2$ . Finally, the forbidden colors for  $v_2v_3$  are in  $\{1, 2, 3, w_1, w_2, w_3, r_1, r_2\}$ .  
 224 Therefore, there is an admissible color  $w_4$  for  $v_2v_3$ .

225 For the subcase depicted in Figure 7(b), the arguments are the same as in  
 226 Figure 7(a) except for  $vv_3$ , which has forbidden colors from  $\{1, 2, 3, w_1, r_2\}$  and  
 227 the three colors used in the neighborhood of  $u$ . So there is an admissible color  
 228  $w_2$  for  $vv_3$ .

229 Subcase B.1.2.  $\deg_G(u) = 4$ . We distinguish several cases. In each case  $\Delta(G') \leq$   
 230  $\Delta(G)$  and  $|C'| < |C|$  hold.

231 (1)  $u = y_3$ ,  $u$  is adjacent to neither  $x_1$  nor  $x_3$ , and  $|\{\psi(uw), \psi(uz)\} \cap \{\psi(x_1x_2),$   
 232  $\psi(x_1x_3)\}| \leq 1$ , where  $z$  is the fourth neighbor of  $u$ , as shown in Figure 8(a). With-  
 233 out loss of generality, assume that  $\psi(uz) \notin \{\psi(x_1x_2), \psi(x_1x_3)\}$ . Let  $\phi(v_1v_2) =$   
 234  $\psi(uz) = 3$  and  $\phi(v_2v_3) = \psi(uw) = 4$ , as indicated in Figure 8(a). Note,  $t_1, t_2 \neq 3$ .  
 235 The forbidden colors for  $vv_1$  are in  $\{1, 2, 3, 4, 5, 6, t_1, t_2\}$ . So there is an admis-  
 236 sible color for  $w_1$ . Next, the forbidden colors for  $w_2$  are in  $\{1, 2, 3, 4, 5, 6, w_1, s\}$ .  
 237 Again, there is an admissible color for  $w_2$ . The forbidden colors for  $w_3$  are in  
 238  $\{1, 2, 3, 4, 5, 6, w_1, w_2\}$ , so there is an admissible color for  $w_3$ .

239 (2)  $u = y_3$ ,  $u$  is adjacent to neither  $x_1$  nor  $x_3$ , and  $\{\psi(uw), \psi(uz)\} =$   
 240  $\{\psi(x_1x_2), \psi(x_1x_3)\}$ , where  $z$  is the fourth neighbor of  $u$ . Without loss of gen-  
 241 erality, we assume that  $\psi(x_1x_2) = \psi(uw) = 5$  and  $\psi(x_1x_3) = \psi(uz) = 7$ . Let  
 242  $\psi(uw) = 3$ ,  $\phi(v_1v_2) = \psi(uy_1) = 4$ ,  $\phi(v_2v_3) = 5$ , and  $\phi(vv_2) = \psi(y_1y_2) = 6$ , as  
 243 indicated in Figure 8(b). Clearly, the remaining edges  $vv_1$  and  $vv_3$  can be colored  
 244 by any two colors not in the set  $\{1, 2, 3, \dots, 7\}$ .

245 (3)  $u = y_3$  and  $u = x_3$  (that is,  $u$  is adjacent to both  $y_1$  and  $x_1$ ). Let  
 246  $\phi(v_1v_2) = \psi(uy_1) = 3$ ,  $\phi(v_2v_3) = \psi(uw) = 4$  and  $\phi(vv_2) = \psi(y_1y_2) = 5$  as  
 247 indicated in Figure 8(c). We find admissible colors  $w_1$  and  $w_2$ . The forbidden  
 248 colors for  $vv_1$  are in  $\{1, 2, 3, 4, 5, 6, 7, t_1\}$ . Hence, there is an admissible color  $w_1$   
 249 for  $vv_1$ . Then the forbidden colors for  $vv_3$  are in  $\{1, 2, 3, 4, 5, 6, 7, w_1\}$ . Thus,




Figure 8.: Subcase B.1.2.

250 there is an admissible color  $w_2$  for  $vv_3$ .

251 (4)  $u$  is adjacent to  $y_3$ ,  $u = x_3$ , and  $\deg_G(y_3) = 3$ . (Symmetrically,  $u$  is  
252 adjacent to  $x_3$ ,  $u = y_3$ , and  $\deg_G(x_3) = 3$ .) Take  $P = y_1, y_3, u, w, u_4, \dots, u_l$  as  
253 a longest path, and such a graph was discussed in Subcase A.3 (see Figure 6(b)  
254 where the positions of  $y_3$  and  $v$  are switched).

255 (5)  $u$  is adjacent to  $y_3$ ,  $u = x_3$ , and  $\deg_G(y_3) = 4$ . Let  $z$  be the fourth neigh-  
256 bor of  $y_3$ . (Symmetrically,  $u$  is adjacent to  $x_3$ ,  $u = y_3$ , and  $\deg_G(x_3) = 4$ .) The  
257 reduction  $G'$  and partial labels are shown in Figure 8(d). The forbidden colors for  
258  $vv_2$  are in  $\{1, 2, 3, 4, 5, 6, 7\}$ . Hence, there is an admissible color  $w_1$  for  $vv_2$ . The  
259 forbidden colors for  $y_2z$  are in  $\{1, 2, 3, 4, 5, 7, s_1, s_2\}$ . Thus, there is an admissi-  
260 ble color  $w_2$  for  $y_2z$ . The forbidden colors for  $y_1y_3$  are from  $\{1, 2, 3, 4, 5, 6, 7, w_2\}$ ,  
261 leaving an admissible color  $w_3$  for  $y_1y_3$ .

262 (6)  $u$  is adjacent to both  $x_3$  and  $y_3$ , and  $\deg_G(x_3) = 3$  or  $\deg_G(y_3) = 3$ . Say  
263  $\deg_G(x_3) = 3$  (the other case is symmetric). Then take  $P = x_1, x_3, u, w, u_4, \dots, u_l$   
264 as a longest path, and such case has been discussed in Case A (see Figure 6).

265 (7)  $u$  is adjacent to both  $x_3$  and  $y_3$ , and  $\deg_G(x_3) = \deg_G(y_3) = 4$ . The  
266 reduction  $G'$  and partial labels are indicated in Figure 8(e). Since  $\deg_G(u_{l-1}) \geq$   
267  $\deg_G(v) = 4$ , we have  $\Delta(G') = \Delta(G)$ . The forbidden colors for  $y_2z_1$  are from  
268  $\{1, 2, 3, 5, 6, 7, s_1, s_2\}$ . Hence, there is an admissible color  $w_1$  for  $y_2z_1$ . The for-  
269 bidden colors for  $y_2y_3$  are in  $\{1, 2, 3, 4, 5, 6, 7, w_1\}$ . Thus, there is an admissible  
270 color  $w_2$  for  $y_2y_3$ . The forbidden colors for  $vv_2$  are from  $\{1, 2, 3, 4, 5, 6, 7\}$ . So  
271 there is an admissible color  $w_3$  for  $vv_2$ .

272 (8)  $u$  is adjacent to  $y_3$ , but not  $x_1$  nor  $x_3$ . Then  $u$  must have another neighbor,  
273 say  $z$ , besides  $y_3$ , that is a leaf or distance one away from the adjoining cycle  $C$ .  
274 The position of  $z$  will be similar to the one on Figure 8(b) (where  $z$  might be on  
275 the cycle). We then consider the longest path  $P^* = y_1y_3u \dots u_l$ , which falls in  
276 one of the cases discussed earlier.

277 **B.2.  $\Delta(G) \geq 5$ .** Obtain the reduction  $G'$  by adding two new edges  $vx_1$  and  $vy_1$   
278 to the induced subgraph of  $G$  on the vertex set  $V(G) \setminus \{v_1, v_2, \dots, v_k\}$ ,  $k \geq 3$ ,  
279 as shown in Figure 9. Since  $\deg_G(u_{l-1}) \geq \deg_G(v)$ , we have  $\Delta(G) = \Delta(G')$ ,  
280 and  $|C'| < |C|$  holds. Without loss of generality, let  $\phi(v_1x_1) = \psi(vx_1) = 1$  and  
281  $\phi(v_ky_1) = \psi(vy_1) = 2$ .

282 For  $u = y_3$  (or  $u$  is adjacent to  $y_3$ , respectively), let  $\phi(vv_2) = \psi(y_1y_2) = 3$   
283 ( $\phi(vv_2) = \psi(y_1y_3) = 3$ , respectively) as indicated in Figure 9(a) (Figure 9(b),  
284 respectively). If  $\deg_G(v) = 4$ , then the coloring scheme is the same as the ones  
285 used in Subcase B.1.1.

286 Thus we assume  $\deg_G(v) \geq 5$ . We proceed to color the remaining edges,  $vv_1$ ,  
287  $vv_3, \dots, vv_k$  and  $v_jv_{j+1}$ , for  $j = 1, 2, \dots, k-1$ .

288 For  $u = y_3$  (see Figure 9(a)), the forbidden colors for  $vv_1$  are  $\{1, 2, 3, t_1, t_2\}$   
289 and colors used in the neighborhood of  $u$ . So there are at most  $\Delta(G) + 5 \leq 2\Delta(G)$   
290 forbidden colors. Hence, there exists an admissible color for  $vv_1$ . Next we color




Figure 9.: Case B.2.

291  $vv_k$ , which has forbidden colors  $\{1, 2, 3, \phi(vv_1)\}$  and the labels used for edges  
 292 incident to  $u$ . Again, there is an admissible color for  $vv_k$ . For  $i = 3, 4, \dots, k - 1$ ,  
 293 we color  $vv_i$  one after another. By direct calculation, the number of forbidden  
 294 colors for  $vv_i$  is at most  $\deg_G(u) + \deg_G(v)$ . Hence, we can color all  $vv_i$  by  
 295 admissible colors.

296 Next we color  $v_1v_2$ , which has forbidden colors  $\{1, t_1, t_2\}$  and colors used  
 297 in the neighborhood of  $v$ . Hence there is an admissible color for  $v_1v_2$ . Next  
 298 we sequentially color  $v_jv_{j+1}$  for  $j = 2, 3, \dots, k - 2$ . Using the assumption that  
 299  $\Delta(G) \geq 5$ , one can easily verify that there exists an admissible color at each step.  
 300 Finally, the forbidden colors for  $v_{k-1}v_k$  are  $\{2, s, \phi(v_{k-2}v_{k-1}), \phi(v_{k-3}v_{k-2})\}$  and  
 301 the labels used in the neighborhood of  $v$ . Thus we can find an admissible color  
 302 for  $v_{k-1}v_k$ .

303 For the case that  $u$  is adjacent to  $y_3$ , the argument is the same except for  
 304 the edge  $vv_k$ , which has forbidden colors from  $\{1, 2, 3, s, \phi(vv_1)\}$  and the labels  
 305 used by the edges incident to  $u$ . As  $\Delta(G) \geq 5$ , we can find an admissible color  
 306 for  $vv_k$ . This completes the proof of Theorem 4. ■

307

### 3. PROOF OF THEOREM 5

308 Let  $G = T \cup C$  be a Halin graph with  $\Delta(G) = 4$ , and let  $G$  be different from  
 309 a wheel. By Theorem 4, if  $\chi'_s(T) = 7$ , then  $\chi'_s(G) \leq \chi'_s(T) + 2$ . So Theorem 5  
 310 holds. Thus we assume  $\chi'_s(T) = 6$ . That is, every vertex of degree 4 is adjacent  
 311 to vertices of degree 3 only. Similarly to the previous section we proceed by  
 312 induction on  $|C|$ , the length of  $C$ . If  $|C| = 4$ , then  $G = W_4$  which contradicts  
 313 the assumption. If  $|C| = 5$ , then  $T = D_{3,4}$  is a double star. The result follows  
 314 by Lemma 6. If  $|C| = 6$ , the only three possible graphs are in Figure 2(a), 2(b),

315 and 2(c). So the result follows.

316 Similarly to the proof of Theorem 4, we consider a reduction  $G' = T' \cup C'$  of  
 317  $G$  with characteristic tree  $T'$  and adjoint cycle  $C'$ . If  $\Delta(G') = 4$  and  $G'$  is not a  
 318 wheel, then  $\chi'_s(G') \leq \chi'_s(T') + 2 \leq \chi'_s(T) + 2$  follows by the induction hypothesis  
 319 since  $|C'| < |C|$ . If  $G' = W_4$  or if  $G'$  is a cubic Halin graph different from  $Ne_2$ ,  
 320 then  $\chi'_s(G') \leq 8 = \chi'_s(T) + 2$  by Theorem 3, Lemma 6, and Lemma 7. Finally,  
 321 the case when  $G' = Ne_2$  is considered at the end of the proof.

322 Assume  $|C| \geq 7$ . Let  $P = u_0, u_1, \dots, u_l$  be a longest path in  $T$  where  $l$  is  
 323 the length of  $P$ . The result holds if  $T$  is a double star by Lemma 6 (note that  
 324  $b \geq 4$ ). Thus, we assume  $l \geq 4$ . Without loss of generality, we also assume that  
 325  $\deg_G(u_1) \leq \deg_G(u_{l-1})$ .

326 Case A. There exists a longest path  $P$  with both non-leaf ends of degree 4. That  
 327 is,  $\deg_G(u_1) = \deg_G(u_{l-1}) = 4$ . Then  $\deg_G(u_2) = 3$ . Consider the following two  
 328 cases.

329 A.1. In  $T$ ,  $u_2$  has exactly one neighbor that is a leaf.




Figure 10.: Case A.1.

330 The reduction  $G'$  along with proposed colors for some edges are depicted in  
 331 Figure 10. We now find admissible colors  $w_1, w_2, w_3, w_4$ , and  $w_5$ . First we can  
 332 find an admissible color  $w_1$  for  $u_1u_2$  that is different from 1, 2 and the colors used  
 333 in the neighborhood of  $u_3$ . Next, we can find an admissible color  $w_2$  for  $v_1v_2$   
 334 that is not in  $\{1, 2, 3, 4, 5, w_1\}$ . Finally, we find three pairwise distinct admissible  
 335 colors  $w_3, w_4, w_5$ , which are not in  $\{1, 2, 3, w_1, w_2\}$ .

336 A.2. In  $T$ , none of the neighbors of  $u_2$  is a leaf.

337 Without loss of generality, we assume that the colors assigned by  $\psi$  to the  
 338 edges incident to  $u_3$  are 3, 4, 5, and 6 (if  $u_3$  has degree 3, then we only use colors 3,  
 339 4, and 5, and ignore the respective edge labeled by 6 in Figure 11). Consider two  
 340 possibilities. For the graph depicted in each Figure 11(a) and 11(b) we obtain the  
 341 reduction  $G'$  and complete the labeling  $\phi$  by using only eight colors, respectively.




Figure 11.: Case A.2.

342 Case B. Every longest path  $P$  has  $\deg_G(u_1) = 3$ . That is, at least one non-leaf  
 343 end has degree 3.

344 B.1.  $\deg_G(u_2) = 3$ .

345 Subcase B.1.1. In  $T$ ,  $u_2$  has exactly one neighbor that is a leaf.




Figure 12.: Subcase B.1.1.

346 The reduction  $G'$  along with proposed colors for some edges are depicted in  
 347 Figure 12. Note if  $u_3$  has degree 3, we simply ignore the edge labeled by  $t_3$  in  
 348 Figure 12. We color  $u_1u_2$  by a color  $w_1$  not from  $\{1, 2, 3, t_1, t_2, t_3\}$ . Next, color  
 349  $v_1v_2$  by a color  $w_2$  not from  $\{1, 2, 3, 4, 5, w_1\}$ . Finally, color  $u_1v_1$  by an admissible  
 350 color  $w_3$  not in  $\{1, 2, 3, w_1, w_2\}$ .

351 Subcase B.1.2. In  $T$ , none of the neighbors of  $u_2$  is a leaf. Then  $u_2$  has two neigh-  
 352 bors, denoted as  $u_1$  and  $v_4$ , that are distance one away from the adjoining cycle  
 353  $C$ . First consider the case that  $v_4$  has degree 4. Then by our assumption of Case  
 354 B, the degree of the other non-leaf end of the path  $P$  must have degree 3. We  
 355 consider the reverse order of  $P$ , denoted as  $P^*$ , as our longest path. That is,  
 356  $P^* = u_l, u_{l-1}, u_{l-2}, \dots, u_1, u_0$ , where  $\deg_G(u_{l-1}) = 3$ . If  $P^*$  falls again in Subcase

357 B.1.2,  $\deg_G(u_{l-2}) = 3$  and none of the neighbors of  $u_{l-2}$  is a leaf, then by the  
 358 assumption of Case B, every non-leaf neighbor of  $v_{l-2}$  that is distance two away  
 359 from the adjoining cycle  $C$  must be degree 3 (for otherwise, there is a longest  
 360 path with both non-leaf ends of degree 4, which was discussed in Case A).

361 Therefore, we only need to consider the case that  $\deg_G(v_4) = 3$ , which is  
 362 shown in Figure 13, where the reduction  $G'$  and partial labels are indicated.




Figure 13.: The second possibility of Subcase B.1.2.

363 We shall find colors for the remaining edges. First, color  $v_3v_4$  and  $v_1v_2$  by  
 364 two admissible colors  $w_1$  and  $w_2$  different from  $\{1, 2, 3, 4, 5\}$ . Next, color  $v_2v_4$  and  
 365  $v_1v_2$  by two admissible colors  $w_3$  and  $w_4$  not from  $\{1, 2, 3, w_1, w_2\}$ , and assign  $u_1v_1$   
 366 the color  $w_5 = w_1$ . Finally color  $u_0u_1$  by an admissible color  $w_6$  different from  
 367  $\{1, 2, 3, w_4, w_5, t_1, t_2\}$ . Since we have 8 colors, this can be accomplished.

368 B.2.  $\deg_G(u_2) = 4$ . Then  $\deg_G(u_3) = 3$ .

369 Subcase B.2.1. In  $T$ ,  $u_2$  has exactly two neighbors that are leaves.

370 Consider possible situations depicted in Figure 14. Figure 14(a) shows the  
 371 situation that the two leaves are adjacent on  $C$ . We color  $v_2v_3$  by a color  $w_1$  not  
 372 from the set  $\{1, 2, 3, 4, 5, s_1, s_2\}$ . Next, color  $u_2v_2$  and  $u_1u_2$  by two colors  $w_2$  and  
 373  $w_3$  not in  $\{1, 2, 3, 4, 5, w_1\}$ .

374 Now assume the two leaves are not adjacent on  $C$ . The length of a longest  
 375 path from  $u_3$  to the adjoint cycle  $C$  on one side of  $v_1$  is at most three, as  $P$  is a  
 376 longest path. Suppose the length is one. Then there is only one possibility which  
 377 is shown in Figure 14(b). Color  $u_2v_4$  by a color  $w_1$  not in  $\{1, 2, 3, 4, 5, t_1, t_2\}$ .  
 378 Color  $u_2v_2$  by a color  $w_2$  not in  $\{1, 2, 3, 4, 5, 6, w_1\}$ . Finally, color  $u_1u_2$  by a color  
 379  $w_3$  not in  $\{1, 2, 3, 4, 5, w_1, w_2\}$ .

380 If there is a path of length two from  $u_3$  to the adjoint cycle  $C$ , then there  
 381 are two possibilities as shown in Figure 14(c) and Figure 14(d). Assume that the  
 382 colors used in the neighborhood of  $u_4$  are from the set  $\{3, 4, 5, 8\}$ . We directly  
 383 color the remaining edges as depicted on those two figures.




Figure 14.: Five possibilities of Subcase B.2.1.



384 Assume there is a path of length three from  $u_3$  to the adjoint cycle  $C$  which  
 385 intersects  $P$  only at  $u_3$ . Let  $u_3, v_2, v_1, v_0$  be such a path from  $u_3$  to  $C$ . Then there  
 386 is another longest path in  $T$ ,  $P' : u_l, u_{l-1}, \dots, u_3, v_2, v_1, v_0$ . Assume  $\deg_G(v_1) = 4$ .  
 387 By our assumption that every longest path has at least one non-leaf end of degree  
 388 3, it must be that  $\deg_G(u_{l-1}) = 3$ . We then consider  $P^*$ , the reverse ordering  
 389 of  $P$ , namely,  $P^* = u_l, u_{l-1}, \dots, u_1, u_0$ . Observe that the same situation will not  
 390 occur to  $P^*$ , since if  $\deg_G(u_{l-2}) = 4$ ,  $\deg_G(u_{l-3}) = 3$ , there is a path of length  
 391 three from  $u_{l-3}$  to  $C$  (denoted as  $u_{l-3}, v'_2, v'_1, v'_0$ ), and  $\deg_G(v'_1) = 4$ , then we  
 392 obtain a longest path  $v'_0, v'_1, v'_2, u_{l-3}, \dots, u_0$  with both non-leaf ends of degree 4,  
 393 which has been discussed in Case A.

394 Thus, assume  $\deg_G(v_1) = 3$ . By symmetry of considering  $P$  and  $P'$ , the only  
 395 possibility is drawn in Figure 14(e), in which an extended strong edge-coloring is  
 396 shown using 8 colors.

397 Subcase B.2.2. In  $T$ ,  $u_2$  has exactly one neighbor that is a leaf.




Figure 15.: Two possibilities of Subcase B.2.2.

398 There are two possible situations as shown in Figure 15. In Figure 15(a),  
 399 a strong edge-coloring is given on the extended edges of  $G'$ . In Figure 15(b),  
 400 we color the edges by the following sequence: Color the two edges labeled as  $w_1$   
 401 by an admissible color not from  $\{1, 2, 3, t_1, t_2\}$ . Color the two edges labeled as  
 402  $w_2$  by an admissible color not from  $\{1, 2, 3, w_1, s_1, s_2\}$ . Color the edge labeled  
 403 as  $w_3$  by an admissible color not from  $\{1, 2, 3, 4, 5, w_1, w_2\}$ . Finally, color the  
 404 remaining two edges labeled as  $w_4$  and  $w_5$  by two different admissible colors not  
 405 from  $\{1, 2, 3, w_1, w_2, w_3\}$ .

406 Subcase B.2.3. In  $T$ , none of the neighbors of  $u_2$  is a leaf. The reduction  $G'$  and  
 407 the completion of  $\phi$  using eight colors are demonstrated in Figure 16. This  
 408 completes all cases.

409 We now discuss the situation that the reduction graph  $G'$  is  $Ne_2$ . Notice that  
 410 this does not occur in Case A. For Subcase B.1.1, if  $G' = Ne_2$ , then  $G$  is a cubic  
 411 graph, contradicting our assumption that  $\chi'_s(T) = 6$ . Similarly, for the second




Figure 16.: Subcase B.2.3.

412 possibility in Subcase B.1.2,  $G'$  is not  $Ne_2$ .

413 These leave a total of fourteen possible situations from the first possibility  
 414 (Figure 11(b)) of Subcase B.1.2, as well as Subcases B.2.1, B.2.2 and B.2.3,  
 415 when the reduction graph  $G'$  is  $Ne_2$ . These fourteen situations are depicted in  
 416 Figure 17, where a strong edge coloring using at most eight colors is given in each  
 417 situation. This completes the proof of Theorem 5.

418 **Acknowledgment.** The authors are grateful of the three anonymous referees for  
 419 careful reading of the manuscript and for helpful constructive comments. Part  
 420 of the work was done when Daphne Liu was visiting the National Center for  
 421 Theoretical Sciences, Taiwan. She is very much thankful for the Center's great  
 422 hospitality.

423

## REFERENCES

- 424 [1] L. D. Andersen, *The strong chromatic index of a cubic graph is at most 10*,  
 425 *Discrete Math.* **108** (1992) 231–252.  
 426 doi:10.1016/0012-365X(92)90678-9
- 427 [2] J. Bensmail, A. Harutyunyan, H. Hocquard and P. Valicov, *Strong edge-*  
 428 *coloring of sparse planar graphs*, *Discrete Appl. Math.* **179** (2014) 229–234.  
 429 doi:10.1016/j.dam.2014.07.006
- 430 [3] O. Borodin and A. Ivanova, *Precise upper bound for the strong edge chro-*  
 431 *matic number of sparse planar graphs*, *Discuss. Math. Graph Theory* **33**  
 432 (2013) 759–770.
- 433 [4] V. Borozan, L. Montero and N. Narayannan, *Further results on strong edge-*




Figure 17.: Fourteen special graphs.

- 434 *colourings in outerplanar graphs*, Australas. J. Combin. **62** (2015) 35–44.  
435 doi:10.7151/dmgt.1708
- 436 [5] R. A. Brualdi and J. Q. Massey, *Incidence and strong edge colorings of*  
437 *graphs*, Discrete Math. **122** (1993) 51–58.  
438 doi:10.1016/0012-365X(93)90286-3
- 439 [6] H. Bruhn and F. Joos, *A stronger bound for the strong chromatic index*,  
440 Electron. Notes in Discrete Math. **49** (2015) 277–284.  
441 doi:10.1016/j.endm.2015.06.038
- 442 [7] K. Cameron, *Induced matchings*, Discrete Appl. Math. **24** (1989) 97–102.  
443 doi:10.1016/0166-218X(92)90275-F
- 444 [8] G. J. Chang and G.-H. Duh, *On the precise value of the strong chromatic-*  
445 *index of a planar graph with a large girth*.  
446 arXiv: 1508.03052.
- 447 [9] G. J. Chang and D. Liu, *Strong edge coloring for cubic Halin graphs*, Discrete  
448 Math. **312** (2012) 1468–1475.  
449 doi:10.1016/j.disc.2012.01.014
- 450 [10] G. J. Chang, M. Montassier, A. Pêcher and A. Raspaud, *Strong chromatic*  
451 *index of planar graphs with large girth*, Discuss. Math. Graph Theory **34**  
452 (2014) 723–733.  
453 doi:10.7151/dmgt.1763
- 454 [11] D. Cranston, *Strong edge-coloring graphs with maximum degree 4 using 22*  
455 *colors*, Discrete Math. **306** (2006) 2772–2778.  
456 doi:10.1016/j.disc.2006.03.053
- 457 [12] P. DeOrsey, J. Diemunsch, M. Ferrara, N. Graber, S. G. Hartke, S. Ja-  
458 hanbekam, B. Lidicky, L. Nelsen, D. Stolee and E. Sullivan, *On the strong*  
459 *chromatic index of sparse graphs*,  
460 arXiv: 1508.03515.
- 461 [13] P. Erdős, *Problems and results in combinatorial analysis and graph theory*,  
462 Discrete Math. **72** (1988) 81–92.  
463 doi:10.1016/0012-365X(88)90196-3
- 464 [14] P. Erdős, J. Nešetřil, *Problem*, in: Irregularities of Partitions, G. Halász and  
465 V. T. Sós (Eds.) (Springer, Berlin, 1989) 162–163.
- 466 [15] R. J. Faudree, R. H. Schelp, A. Gyárfás and Zs. Tuza, *The strong chromatic*  
467 *index of graphs*, Ars Combin. **29B** (1990) 205–211.

- 468 [16] J. L. Fouquet and J. Jolivet, *Strong edge-coloring of graphs and applications*  
469 *to multi- $k$ -gons*, Ars Combin. **16A** (1983) 141–150.
- 470 [17] J. L. Fouquet and J. Jolivet, *Strong edge-coloring of cubic planar graphs*,  
471 Progress in Graph Theory (Academic Press, Toronto, 1984) 247–264.
- 472 [18] M. C. Golumbic and M. Lewenstein, *New results on induced matchings*,  
473 Discrete Appl. Math. **101** (2000) 157–165.  
474 doi:10.1016/S0166-218X(99)00194-8
- 475 [19] H. Hocquard, M. Montassier, A. Raspaud and P. Valicov, *On strong edge-*  
476 *colouring of subcubic graphs*, Discrete Appl. Math. **161** (2013) 2467–2479.  
477 doi:10.1016/j.dam.2013.05.021
- 478 [20] H. Hocquard, P. Ochem and P. Valicov, *Strong edge-colouring and induced*  
479 *matchings*, Inform. Process. Lett. **113** (2013) 836–843.  
480 doi:10.1016/j.ipl.2013.07.026
- 481 [21] P. Horák, *The strong chromatic index of graphs with maximum degree*  
482 *four*, Contemporary Methods in Graph Theory (Bibliographisches Inst.,  
483 Mannheim, 1990) 399–403.
- 484 [22] P. Horák, H. Qing and W. T. Trotter, *Induced matchings in cubic graphs*,  
485 J. Graph Theory **17** (1993) 151–160.  
486 doi:10.1002/jgt.3190170204
- 487 [23] D. Hudák, B. Lužar, R. Soták and R. Škrekovski, *Strong edge coloring of*  
488 *planar graphs*, Discrete Math. **324** (2014) 41–49.  
489 doi:10.1016/j.disc.2014.02.002
- 490 [24] H.-H. Lai, K.-W. Lih and P.-Y. Tsai, *The strong chromatic index of Halin*  
491 *graphs*, Discrete Math. **312** (2012) 1536–1541.  
492 doi:10.1016/j.disc.2011.09.016
- 493 [25] K.-W. Lih and D. D.-F. Liu, *On the strong chromatic index of cubic Halin*  
494 *graphs*, Appl. Math. Lett. **25** (2012) 898–901.  
495 doi:10.1016/j.aml.2011.10.046
- 496 [26] M. Mahdian, *On the computational complexity of strong edge coloring*, Dis-  
497 crete Appl. Math. **118** (2002) 239–248.  
498 doi:10.1016/S0166-218X(01)00237-2
- 499 [27] M. Maydanskiy, *The incidence coloring conjecture for graphs of maximum*  
500 *degree 3*, Discrete Math. **292** (2005) 131–141.  
501 doi:10.1016/j.disc.2005.02.003

- 502 [28] M. Molloy and B. Reed, *A bound on the strong chromatic index of a graph*,  
503 J. Combin. Theory Ser. B **69** (1997) 103–109.  
504 doi:10.1006/jctb.1997.1724
- 505 [29] M. R. Salavatipour, *A polynomial time algorithm for strong edge coloring of*  
506 *partial  $k$ -trees*, Discrete Appl. Math. **143** (2004) 285–291.  
507 doi:10.1016/j.dam.2004.03.001
- 508 [30] W. C. Shiu, P. C. B. Lam and W. K. Tam, *On strong chromatic index of*  
509 *Halin graphs*, J. Combin. Math. Combin. Comput. **57** (2006) 211–222.
- 510 [31] W. C. Shiu and W. K. Tam, *The strong chromatic index of complete cubic*  
511 *Halin graphs*, Appl. Math. Lett. **22** (2009) 754–758.  
512 doi:10.1016/j.aml.2008.08.019
- 513 [32] J. Wu and W. Lin, *The strong chromatic index of a class of graphs*, Discrete  
514 Math. **308** (2008) 6254–6262.  
515 doi:10.1016/j.disc.2007.11.051