

# Distance graphs with missing multiples in the distance sets

Daphne Der-Fen Liu \*

Department of Mathematics and Computer Science  
California State University, Los Angeles  
Los Angeles, CA 90032, USA  
Email: dliu@calstatela.edu

Xuding Zhu †

Department of Applied Mathematics  
National Sun Yat-sen University  
Kaoshing, Taiwan 80424  
Email: zhu@ibm18.math.nsysu.edu.tw

July 13, 1997 (Revised October 2, 1998)

## Abstract

Given positive integers  $m, k$  and  $s$  with  $m > ks$ , let  $D_{m,k,s}$  represent the set  $\{1, 2, \dots, m\} - \{k, 2k, \dots, sk\}$ . The distance graph  $G(Z, D_{m,k,s})$  has as vertex set all integers  $Z$  and edges connecting  $i$  and  $j$  whenever  $|i - j| \in D_{m,k,s}$ . The chromatic number and the fractional chromatic number of  $G(Z, D_{m,k,s})$  are denoted by  $\chi(Z, D_{m,k,s})$  and  $\chi_f(Z, D_{m,k,s})$ , respectively. For  $s = 1$ ,  $\chi(Z, D_{m,k,1})$  was studied by Eggleton, Erdős and Skilton [6], Kemnitz and Kolberg [12], and Liu [13], and was solved lately by Chang, Liu and Zhu [2] who also determined  $\chi_f(Z, D_{m,k,1})$  for any  $m$  and  $k$ . This article extends the study of  $\chi(Z, D_{m,k,s})$  and  $\chi_f(Z, D_{m,k,s})$  to general values of  $s$ . We prove  $\chi_f(Z, D_{m,k,s}) = \chi(Z, D_{m,k,s}) = k$  if  $m < (s + 1)k$ ; and  $\chi_f(Z, D_{m,k,s}) = (m + sk + 1)/(s + 1)$  otherwise. The latter result provides a good lower bound for  $\chi(Z, D_{m,k,s})$ . A general upper bound for  $\chi(Z, D_{m,k,s})$  is found. We prove the upper bound can be improved to  $\lceil (m + sk + 1)/(s + 1) \rceil + 1$  for some values of  $m, k$  and  $s$ . In particular, when  $s + 1$  is prime,  $\chi(Z, D_{m,k,s})$  is either  $\lceil (m + sk + 1)/(s + 1) \rceil$  or  $\lceil (m + sk + 1)/(s + 1) \rceil + 1$ . By using a special coloring method called the pre-coloring method, many distance graphs  $G(Z, D_{m,k,s})$  are classified into

---

\*Supported in part by the National Science Foundation under grant DMS 9805945.

†Supported in part by the National Science Council under grant NSC87-2115-M110-004.

these two possible values of  $\chi(Z, D_{m,k,s})$ . Moreover, complete solutions of  $\chi(Z, D_{m,k,s})$  for several families are determined including the case  $s = 1$  (solved in [2]), the case  $s = 2$ , the case  $(k, s + 1) = 1$ , and the case that  $k$  is a power of a prime.

**Keywords.** Distance graph, chromatic number, fractional chromatic number, pre-coloring method.

## 1 Introduction

Given a set  $D$  of positive integers, the *distance graph*  $G(Z, D)$  has all integers as vertices; and two vertices are adjacent if and only if their difference falls within  $D$ , that is, the vertex set is  $Z$  and the edge set is  $\{uv : |u - v| \in D\}$ . We call  $D$  the *distance set*. The chromatic number of  $G(Z, D)$  is denoted by  $\chi(Z, D)$ .

For different types of distance sets  $D$ , the problem of determining  $\chi(Z, D)$  has been studied extensively. (See [2, 3, 4, 6, 7, 8, 9, 12, 16, 15, 17].) For instance, suppose  $D$  is a subset of prime numbers and  $\{2, 3\} \subseteq D$ , Eggleton, Erdős and Skilton [9] proved that  $\chi(Z, D)$  is either 3 or 4. The problem of classifying  $G(Z, D)$  with distance sets  $D$  of primes into chromatic number 3 or 4 was studied by Eggleton, Erdős and Skilton [9], and by Voigt and Walther [16]. However, a complete classification is not obtained yet.

If  $D$  has only one element, it is trivial that  $\chi(Z, D) = 2$ . When  $D$  has two elements, it is known that  $\chi(Z, D) = 3$  if the two integers in  $D$  are of different parity, and  $\chi(Z, D) = 2$  otherwise (assuming that  $\gcd D = 1$ ). The case if  $D$  has three elements, which is much more complicated, has been studied by Chen, Chang, and Huang [3], and by Voigt [15], and was solved lately by Zhu [17].

A *fractional coloring* of a graph  $G$  is a mapping  $h$  from  $\mathcal{I}(G)$ , the set of all independent sets of  $G$ , to the interval  $[0, 1]$  such that  $\sum_{I \in \mathcal{I}(G), x \in I} h(I) \geq 1$  for each vertex  $x$  of  $G$ . The *fractional chromatic number*  $\chi_f(G)$  of  $G$  is the infimum of the value  $\sum_{I \in \mathcal{I}(G)} h(I)$  of a fractional coloring  $h$  of  $G$ . The fractional chromatic number of

a distance graph  $G(Z, D)$  is denoted by  $\chi_f(Z, D)$ .

For any graph  $G$ , it is well-known and easy to verify that

$$\max\{\omega(G), \frac{|V(G)|}{\alpha(G)}\} \leq \chi_f(G) \leq \chi(G), \quad (*)$$

where  $\omega(G)$  is the size (number of vertices) of a maximum complete graph, and  $\alpha(G)$  is the size of a maximum independent set in  $G$ . (See Chapter 3 of [14].)

Given integers  $m, k$  and  $s$  with  $m > ks$ , let  $D_{m,k,s}$  denote the distance set  $D_{m,k,s} = \{1, 2, 3, \dots, m\} - \{k, 2k, 3k, \dots, sk\}$ . This article studies the chromatic number and the fractional chromatic number of  $G(Z, D_{m,k,s})$ . If  $s = 1$ , the chromatic number of  $G(Z, D_{m,k,1})$  was first studied by Eggleton, Erdős and Skilton [6] who determined  $\chi(Z, D_{m,k,1})$  completely for  $k = 1$ , and partially for  $k = 2$ . The same results for the case  $k = 1$  were also obtained in [12] by a different approach. For the cases that  $k$  is an odd number,  $k = 2$  and  $k = 4$ ,  $\chi(Z, D_{m,k,1})$  were determined in [13]. Recently, the exact values of  $\chi_f(Z, D_{m,k,1})$  and  $\chi(Z, D_{m,k,1})$  for all  $m$  and  $k$  were settled in [2]. We extend the study to general values of  $s$ .

Note that it becomes an easy case if  $m < (s + 1)k$ . Define a coloring  $f$  of  $G(Z, D_{m,k,s})$  by: For any  $x \in Z$ ,  $f(x) = x \bmod k$ . Since  $D_{m,k,s}$  contains no multiples of  $k$ ,  $f$  is a proper coloring. Thus,  $\chi(Z, D_{m,k,s}) \leq k$ . As any consecutive  $k$  vertices in  $G(Z, D_{m,k,s})$  form a complete graph, by (\*),  $\chi_f(Z, D_{m,k,s}) \geq k$ . This implies  $\chi(Z, D_{m,k,s}) = \chi_f(Z, D_{m,k,s}) = k$ , if  $m < (s + 1)k$ . Therefore, throughout the article, we assume  $m \geq (s + 1)k$ .

Section 2 determines the fractional chromatic number of  $G(Z, D_{m,k,s})$  for all values of  $m, k$  and  $s$  with  $m \geq (s + 1)k$ . This result provides a good lower bound for  $\chi(Z, D_{m,k,s})$ , namely,

$$\lceil (m + sk + 1)/(s + 1) \rceil \leq \chi(Z, D_{m,k,s}), \text{ if } m \geq (s + 1)k. \quad (**)$$

This lower bound will be shown to be sharp for some families of  $G(Z, D_{m,k,s})$  and strict for some others.

Section 3 introduces the pre-coloring method, one of the main tools used in the article. For such a coloring method, we determine when it produces a proper coloring for  $G(Z, D_{m,k,s})$ , and then determine the number of colors used by the produced proper coloring. These characterizations are used intensively in Sections 4 and 5.

Section 4 starts with the result of a general upper bound of  $\chi(Z, D_{m,k,s})$ . For some values of  $m, k$  and  $s$ , we improve the upper bound to  $\lceil (m + sk + 1)/(s + 1) \rceil + 1$ . Combining these results with the lower bound (\*\*\*) mentioned above, the chromatic numbers for many families of  $G(Z, D_{m,k,s})$  are determined.

Section 5 focuses on the study of  $\chi(Z, D_{m,k,s})$  when  $s + 1$  is a prime number. Using the results obtained in earlier sections, we show that when  $s + 1$  is prime,  $\chi(Z, D_{m,k,s})$  is either  $\lceil (m + sk + 1)/(s + 1) \rceil$  or  $\lceil (m + sk + 1)/(s + 1) \rceil + 1$ . For many families of  $G(Z, D_{m,k,s})$ , we classify their chromatic numbers into one of these two values. Moreover, we completely determine the exact values of  $\chi(Z, D_{m,k,s})$  for the following cases: If  $s = 1$  (which was solved recently in [2]); if  $s = 2$ ; if  $(k, s + 1) = 1$ ; and if  $k$  is a power of a prime.

## 2 Lower bounds and fractional chromatic number

In this section, we first determine the fractional chromatic number of  $G(Z, D_{m,k,s})$  for all values of  $m, k$  and  $s$  with  $m \geq (s + 1)k$ . This result immediately leads to (\*\*), a lower bound for  $\chi(Z, D_{m,k,s})$ . Then we prove that in (\*\*), equality holds for some values of  $m, k$  and  $s$ ; while strict inequality holds for some others.

**Theorem 1** *For any given integers  $m, k$  and  $s$  with  $m \geq (s + 1)k$ ,*

$$\chi_f(Z, D_{m,k,s}) = (m + sk + 1)/(s + 1).$$

**Proof.** For any  $i$  with  $0 \leq i \leq m + sk$ , let  $I_i = \{j \in Z : j - i \equiv xk \pmod{m + sk + 1}, 0 \leq x \leq s\}$ . It is straightforward to verify that  $I_i$  is an independent set in

$G(Z, D_{m,k,s})$ . It is also easy to verify that any integer is contained in exactly  $s + 1$  such independent sets. Define a mapping  $h : \mathcal{I}(G(Z, D_{m,k,s})) \rightarrow [0, 1]$  by

$$h(I) = \begin{cases} \frac{1}{s+1}, & \text{if } I = I_i \text{ for } 0 \leq i \leq m + sk; \\ 0, & \text{otherwise.} \end{cases}$$

Then  $h$  is a fractional coloring of  $G(Z, D_{m,k,s})$  which has value  $\frac{m+sk+1}{s+1}$ . Thus,  $\chi_f(Z, D_{m,k,s}) \leq \frac{m+sk+1}{s+1}$ .

To show  $\chi_f(Z, D_{m,k,s}) \geq \frac{m+sk+1}{s+1}$ , let  $G$  be the subgraph of  $G(Z, D_{m,k,s})$  induced by the vertices  $\{0, 1, 2, \dots, m+sk\}$ . Then  $\chi_f(G) \leq \chi_f(Z, D_{m,k,s})$ . It is straightforward to verify that  $\alpha(G) = s + 1$ . Hence, by (\*),  $\chi_f(G) \geq \frac{|V(G)|}{\alpha(G)} = \frac{m+sk+1}{s+1}$ . This completes the proof of Theorem 1. Q.E.D.

Since  $\chi(G)$  is an integer, by (\*), we have  $\lceil \chi_f(G) \rceil \leq \chi(G)$ . Hence, the following is obtained.

**Theorem 2** *For any given integers  $m, k$  and  $s$  with  $m \geq (s + 1)k$ ,*

$$\chi(Z, D_{m,k,s}) \geq \lceil (m + sk + 1)/(s + 1) \rceil.$$

The following result indicates that the lower bound of  $\chi(Z, D_{m,k,s})$  in Theorem 2 is attained by some values of  $m, k$  and  $s$ , but not attained by some others.

**Theorem 3** *Suppose  $m \geq (s + 1)k$ ,  $k = (s + 1)^a k'$  and  $m + sk + 1 = (s + 1)^b m'$ , where both  $k'$  and  $m'$  are not divisible by  $s + 1$ . Then*

$$\chi(Z, D_{m,k,s}) \begin{cases} \geq (m + sk + 1)/(s + 1) + 1, & \text{if } 0 < b \leq a; \\ = (m + sk + 1)/(s + 1), & \text{if } a < b \text{ and } (s + 1, k') = 1. \end{cases}$$

**Proof.** Let  $n = (m + sk + 1)/(s + 1)$ . Because  $b > 0$ ,  $n$  is an integer.

Suppose  $0 < b \leq a$ , we shall show that  $G(Z, D_{m,k,s})$  is not  $n$ -colorable. Assume to the contrary, there exists an  $n$ -coloring  $f$  of  $G(Z, D_{m,k,s})$ .

For any two integers  $i$  and  $j$ , let  $G[i, j]$  be the subgraph of  $G(Z, D_{m,k,s})$  induced by the vertex set  $\{i+1, i+2, \dots, j\}$ . Then for any integer  $i$ , the graph  $G[i, i+m+sk+1]$

has  $m + sk + 1$  vertices and a maximum independent set of size  $s + 1$ . Since  $f$  is an  $(m + sk + 1)/(s + 1)$ -coloring, exactly  $s + 1$  vertices of  $G[i, i + m + sk + 1]$  are colored by the same color. It follows that  $f(i) = f(i + m + sk + 1)$  for any integer  $i$ .

Define a circulant graph  $G$  on the set  $\{0, 1, \dots, m + sk\}$  with generating set  $D_{m,k,s}$ , that is,  $ij$  is an edge of  $G$  if and only if  $(j - i) \bmod (m + sk + 1) \in D_{m,k,s}$  or  $(i - j) \bmod (m + sk + 1) \in D_{m,k,s}$ . The argument in the previous paragraph shows that  $f$  induces a proper  $n$ -coloring of  $G$ . Moreover, each color class consists of  $s + 1$  vertices in  $G$ . It is not difficult to verify that all  $(s + 1)$ -independent sets of  $G$  are of the form  $\{i, i + k, \dots, i + sk\}$ . (Here each number is calculated by modulo  $m + sk + 1$ .)

Let  $d = (k, m + sk + 1)$  and  $u = (m + sk + 1)/d$ . Divide the vertex set of  $G$  into  $d$  subsets of the form  $\{i, i + k, i + 2k, \dots, i + (u - 1)k\} \pmod{m + sk + 1}$ , each of size  $u$ . Then each of these  $d$  subsets is the union of some color classes of size  $s + 1$ , so  $(s + 1)$  divides  $u$ . Therefore  $m + sk + 1$  is a multiple of  $(s + 1)^{a+1}$ , which is impossible since  $b \leq a$ .

Suppose  $a < b$  and  $(s + 1, k') = 1$ , then  $u$  is a multiple of  $s + 1$ . One can easily define a proper  $n$ -coloring  $f$  on  $G$  by using  $u/(s + 1)$  colors to each of the subsets  $\{i, i + k, i + 2k, \dots, i + (u - 1)k\} \pmod{m + sk + 1}$  as defined in the previous paragraph by: the first  $s + 1$  vertices in a subset use one color and the next  $s + 1$  vertices use the next, and continue the process until all vertices are colored. It is easy to check that  $f$  is a proper coloring of  $G$ . Furthermore,  $f$  can be extended to a proper coloring of  $G(Z, D_{m,k,s})$  by letting  $f'(y) = f(x)$ , where  $x = y \bmod (m + sk + 1)$ . Therefore,  $G(Z, D_{m,k,s})$  is  $n$ -colorable. This completes the proof of Theorem 3. Q.E.D.

### 3 The pre-coloring method

This section introduces the main tool to be used in the remaining part of this article, namely, the *pre-coloring method*. A simpler version of this method was originally applied in [2] in determining the chromatic number of  $G(Z, D_{m,k,1})$ . Here we extend

the idea to a more complex version and use it extensively throughout this article.

Before introducing the pre-coloring method, we note another fact. Let  $Z^*$  denote the set of non-negative integers. It is known and easy to verify that for any distance set  $D$ ,  $\chi(Z, D) = \chi(Z^*, D)$ , where  $G(Z^*, D)$  is the subgraph of  $G(Z, D)$  induced by  $Z^*$ . Therefore, to color the graph  $G(Z, D_{m,k,s})$ , it suffices to color the subgraph of  $G(Z, D_{m,k,s})$  induced by  $Z^*$ .

There are two steps in the pre-coloring method. First, we partition the set  $Z^*$  into  $s + 1$  parts by a mapping  $c : Z^* \rightarrow \{0, 1, 2, \dots, s\}$ . Second, for each non-negative integer  $x$ , according to the value of  $c(x)$ , we assign a color to  $x$  by the rule defined as follows.

**Definition 4** *Suppose  $m, k, s$  are positive integers. For a given mapping  $c : Z^* \rightarrow \{0, 1, 2, \dots, s\}$ , define a coloring  $c'$  of  $Z^*$  recursively by:*

$$c'(j) = \begin{cases} j, & \text{if } j < k; \\ c'(j - k), & \text{if } j \geq k \text{ and } c(j) \neq 0; \\ n, & \text{if } j \geq k \text{ and } c(j) = 0, \end{cases}$$

where  $n$  is the smallest non-negative integer (color) not been used in the  $m$  vertices preceding  $j$ , that is,  $n = \min\{t \in Z^* : c'(j - i) \neq t \text{ for } i = 1, 2, \dots, m\}$ .

Note that  $c'$  defined above is uniquely determined by  $c$ . We call  $c$  the *pre-coloring*, and  $c'$  the *coloring induced by  $c$* . For any  $x \in Z^*$ ,  $c(x)$  and  $c'(x)$  are called the *pre-color* and the *color* of  $x$ , respectively.

In order to ensure that the coloring  $c'$  in Definition 4 to be a proper coloring for  $G(Z^*, D_{m,k,s})$  as desired, the pre-coloring  $c$  needs to satisfy certain conditions specified in the following lemma.

**Lemma 5** *Suppose  $c$  is a pre-coloring of  $Z^*$ . If for any integer  $j \geq sk$ ,  $c(j), c(j - k), c(j - 2k), \dots$ , and  $c(j - sk)$  are all distinct, then the induced coloring  $c'$  is a proper coloring for  $G(Z, D_{m,k,s})$ .*

**Proof.** It is enough to show by induction that for any  $j \in Z^*$ ,  $c'(j) \neq c'(x)$  for any neighbor  $x$  of  $j$  and  $x < j$ . If  $j < k$ , or  $j \geq k$  with  $c(j) = 0$ , then this is true by Definition 4.

Now, assume  $j \geq k$  and  $c(j) \neq 0$ . By definition,  $c'(j) = c'(j - k)$ . If  $j - k < x < j$ , then  $x$  is adjacent to  $j - k$ . By the inductive hypotheses,  $c'(x) \neq c'(j - k)$ , so  $c'(x) \neq c'(j)$ . If  $x < j - k$  and  $x$  is adjacent to  $j$ , then either  $x$  is a neighbor of  $j - k$  or  $x = j - (s + 1)k$ . In the former case, according to the inductive hypotheses,  $c'(x) \neq c'(j - k)$ , hence  $c'(x) \neq c'(j)$ . We now consider the case that  $x = j - (s + 1)k$ . Because the pre-colors of  $j, j - k, j - 2k, \dots, j - sk$  are all distinct, exactly one of them is 0. Suppose  $c(j - uk) = 0$  for some  $0 \leq u \leq s$ . Then by Definition 4,  $c'(j - uk)$  is different from the color of any of the  $m$  vertices preceding  $j - uk$ , hence  $c'(j - uk) \neq c'(j - (s + 1)k)$ . Because  $c(j), c(j - k), \dots, c(j - (u - 1)k) \neq 0$ ,  $c'(j) = c'(j - k) = c'(j - 2k) = \dots = c'(j - uk)$ . Therefore,  $c'(j) \neq c'(j - (s + 1)k)$ , *i.e.*,  $c'(j) \neq c'(x)$ . This completes the proof of Lemma 5. Q.E.D.

After getting a necessary condition for the pre-coloring  $c$  to produce a proper coloring  $c'$  for the distance graph  $G(Z^*, D_{m,k,s})$ , the next natural question to ask is *how many* colors are used by  $c'$ . The answer of this question is shown in the following result.

**Lemma 6** *Suppose  $c$  is a pre-coloring and  $c'$  is the induced coloring. Then the number of colors used by  $c'$  is at most  $k + \ell$ , where  $\ell$  is the maximum number of vertices with pre-color 0, among any  $m - k + 1$  consecutive integers greater than  $k$ .*

**Proof.** We prove, by induction on  $j$ , that vertices  $0, 1, 2, \dots, j$  are colored by the pre-coloring method with at most  $k + \ell$  colors. This is trivial when  $j < k$ , or  $j \geq k$  with  $c(j) \neq 0$ .

Now we assume  $j > k$  and  $c(j) = 0$ . It suffices to show that the  $m$  vertices preceding  $j$  use at most  $k + \ell - 1$  colors. For the  $m$  vertices preceding  $j$ , the first  $k$

vertices use at most  $k$  colors. Among the remaining  $m - k$  vertices, only those vertices with pre-color 0 require a new color. Due to the facts that  $c(j) = 0$ , and any set of consecutive  $m - k + 1$  vertices contains at most  $\ell$  vertices of pre-color 0, we conclude that among the remaining  $m - k$  vertices, there are at most  $\ell - 1$  vertices with pre-color 0. Therefore, the total number of colors used by the  $m$  vertices preceding  $j$  is at most  $k + \ell - 1$ , and hence there is a color for the vertex  $j$ . Q.E.D.

Combining Lemmas 5 and 6, we arrive at the following useful conclusion.

**Corollary 7** *Given integers  $m, k$  and  $s$ ,  $\chi(Z, D_{m,k,s}) \leq n$  if there exists a pre-coloring  $c$  such that the following two conditions are satisfied:*

- (1) *for any integer  $j \geq sk$ ,  $c(j), c(j - k), c(j - 2k), \dots, c(j - sk)$  are all distinct,*  
*and*
- (2) *among any consecutive non-negative  $m - k + 1$  integers, there are at most  $n - k$  vertices with pre-color 0.*

Corollary 7 will be used in many of the proofs in the rest of the article. Instead of finding a proper coloring for the distance graph  $G(Z, D_{m,k,s})$  with  $n$  colors, it is enough to present a pre-coloring  $c$  that satisfies (1) and (2) of Corollary 7.

## 4 Upper bounds

This section shows upper bounds of  $\chi(Z, D_{m,k,s})$  for different values of  $m, k$  and  $s$ . Combining these upper bounds with the lower bounds obtained in Section 2 gives the exact value of  $\chi(Z, D_{m,k,s})$  for some families of  $G(Z, D_{m,k,s})$ . In particular, we prove for many different combinations of  $m, k$  and  $s$ ,  $\chi(Z, D_{m,k,s})$  is either  $\lceil (m + sk + 1)/(s + 1) \rceil$  or  $\lceil (m + sk + 1)/(s + 1) \rceil + 1$ .

We start with a general upper bound in the following. For any two integers  $a$  and  $b$ , let  $(a, b)$  denote the greatest common divisor of  $a$  and  $b$ .

**Theorem 8** *Suppose  $m \geq (s + 1)k$  and  $(k, m + sk + 1) = d$ , then  $\chi(Z, D_{m,k,s}) \leq d \lceil (m + sk + 1)/d(s + 1) \rceil$ .*

**Proof.** Define a circulant graph  $G$  on the set  $\{0, 1, \dots, m + sk\}$  with generating set  $D_{m,k,s}$ , that is,  $ij$  is an edge of  $G$  if and only if  $(j - i) \bmod (m + sk + 1) \in D_{m,k,s}$  or  $(i - j) \bmod (m + sk + 1) \in D_{m,k,s}$ . It is easy to verify that any proper coloring  $f$  of  $G$  can be extended to a proper coloring  $f'$  of  $G(Z, D_{m,k,s})$  by letting  $f'(y) = f(x)$ , where  $x = y \bmod (m + sk + 1)$ . Therefore, it is enough to find a proper  $n$ -coloring of  $G$ , where  $n = d \lceil (m + sk + 1)/d(s + 1) \rceil$ .

Let  $u = (m + sk + 1)/d$ . Divide the vertex set of  $G$  into  $d$  subsets such that each subset has  $u$  vertices and is of the form  $\{i, i + k, i + 2k, \dots, i + (u - 1)k\} \pmod{m + sk + 1}$ . Any consecutive  $s + 1$  vertices in a subset are independent, so each subset can be partitioned into  $\lceil u/(s + 1) \rceil = \lceil (m + sk + 1)/d(s + 1) \rceil$  independent sets of size  $s + 1$ , except the last one whose size might be smaller than  $s + 1$ . Therefore the vertex set of  $G$  can be partitioned into  $d \lceil (m + sk + 1)/d(s + 1) \rceil$  independent sets. Hence  $\chi(Z, D_{m,k,s}) \leq d \lceil (m + sk + 1)/d(s + 1) \rceil$ . Q.E.D.

Combining the upper bound above with the lower bound in Theorem 2, the following two results emerge.

**Corollary 9** *Suppose  $m \geq (s + 1)k$  and  $(k, m + sk + 1) = d$ , then*

$$\lceil (m + sk + 1)/(s + 1) \rceil \leq \chi(Z, D_{m,k,s}) \leq d \lceil (m + sk + 1)/d(s + 1) \rceil.$$

**Corollary 10** *If  $m \geq (s + 1)k$  and  $(k, m + sk + 1) = 1$ , then  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s + 1) \rceil$ .*

We note that in Corollary 9, there may exist big gaps between the upper and the lower bounds, depending on the values of  $d = (k, m + sk + 1)$ . However, so far we do not have any example of distance graph  $G(Z, D_{m,k,s})$  with chromatic number

exceeding  $\lceil (m + sk + 1)/(s + 1) \rceil + 1$ . The next theorem provides a better upper bound for some families of  $G(Z, D_{m,k,s})$ .

**Theorem 11** *If  $m \geq (s + 1)k$  and  $s + 1$  is a divisor of  $k$ , then  $\chi(Z, D_{m,k,s}) \leq \lceil (m + sk + 1)/(s + 1) \rceil + 1$ .*

**Proof.** For any  $j \in Z^*$ , we can write  $j$  uniquely in the form  $j = uk + v(s + 1) + w$ , where  $u, v$  and  $w$  are integers such that  $0 \leq v < k/(s + 1)$  and  $0 \leq w \leq s$ . Then define a pre-coloring  $c$  by  $c(j) = u + w \pmod{s + 1}$ . We only need to show that  $c$  satisfies (1) and (2) in Corollary 7, with  $n = \lceil (m + sk + 1)/(s + 1) \rceil + 1$ .

First we show that for any vertex  $j$ , the  $s + 1$  vertices,  $j, j - k, j - 2k, \dots, j - sk$  have distinct pre-colors. Assume  $j = uk + v(s + 1) + w$  with  $0 \leq v < k/(s + 1)$  and  $0 \leq w \leq s$ . Then  $j - ik = (u - i)k + v(s + 1) + w$ ,  $0 \leq i \leq s$ . It follows that  $c(j - ik) = (u - i + w) \pmod{s + 1}$  which give distinct colors for  $0 \leq i \leq s$ .

Next we show that among any consecutive  $m - k + 1$  vertices, there are at most  $n - k = \lceil (m - k + 1)/(s + 1) \rceil + 1$  vertices with pre-color 0. Divide the set of non-negative integers into segments of length  $s + 1$  by  $A_0 = \{0, 1, \dots, s\}, A_1 = \{s + 1, s + 2, \dots, 2s + 1\}, \dots, A_i = \{i(s + 1), i(s + 1) + 1, \dots, (i + 1)(s + 1) - 1\}, \dots$ . Then each segment  $A_i$  contains exactly one vertex of each pre-color. Indeed, it is straightforward to verify that the pre-colors of  $A_i$  are  $\{j, j + 1, \dots, s, 0, 1, \dots, j - 1\}$ , where  $i = uk/(s + 1) + v$ ,  $0 \leq v < k/(s + 1)$  and  $j = u \pmod{s + 1}$ . Any set of consecutive  $m - k + 1$  vertices intersects at most  $\lceil (m - k + 1)/(s + 1) \rceil + 1$  segments, so it contains at most  $\lceil (m - k + 1)/(s + 1) \rceil + 1$  vertices of pre-color 0. This completes the proof. Q.E.D.

The following corollary follows from Theorems 3 and 11.

**Corollary 12** *Suppose  $m \geq (s + 1)k$ ,  $k = (s + 1)^a k'$  and  $m + sk + 1 = (s + 1)^b m'$ , where both  $k'$  and  $m'$  are not divisible by  $s + 1$ . If  $0 < b \leq a$ , then  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s + 1) \rceil + 1$ .*

The next result shows another family of  $G(Z, D_{m,k,s})$  such that the chromatic number reaches the lower bound.

**Theorem 13** *If  $(k, s + 1) = 1$ , then  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s + 1) \rceil$  for all  $m \geq (s + 1)k$ .*

**Proof.** Define a pre-coloring  $c$  by  $c(j) = j \pmod{s + 1}$ . We prove that  $c$  satisfies (1) and (2) of Corollary 7, with  $n = \lceil (m + sk + 1)/(s + 1) \rceil$ .

To show that for any vertex  $j$ ,  $c(j), c(j - k), c(j - 2k), \dots$ , and  $c(j - sk)$  are all distinct, we assume to the contrary that  $c(j - tk) = c(j - t'k)$  for some  $0 \leq t < t' \leq s$ . Then  $j - tk \equiv j - t'k \pmod{s + 1}$ , so  $(t' - t)k \equiv 0 \pmod{s + 1}$ . This is impossible, because  $(k, s + 1) = 1$  and  $0 < t' - t \leq s$ .

Next we show that among any consecutive  $m - k + 1$  vertices, there are at most  $\lceil (m - k + 1)/(s + 1) \rceil$  vertices with pre-color 0. This is trivial, because the vertices of pre-color 0 are those vertices  $j$  for which  $j \equiv 0 \pmod{s + 1}$ , so any two vertices with pre-color 0 are exactly  $s + 1$  vertices apart. This completes the proof. Q.E.D.

## 5 The case $s + 1$ is prime

This section focuses on the study of  $\chi_f(Z, D_{m,k,s})$  when  $s + 1$  is a prime number. If  $s + 1$  is prime, then either  $s + 1$  is a divisor of  $k$  or  $(k, s + 1) = 1$ . Hence by Theorems 11 and 13,  $\chi(Z, D_{m,k,s})$  is either  $\lceil (m + sk + 1)/(s + 1) \rceil$  or  $\lceil (m + sk + 1)/(s + 1) \rceil + 1$ . In this section, assuming  $s + 1$  is prime, we classify the chromatic number for most of the families of the distance graphs  $G(Z, D_{m,k,s})$  into one of those two possible values.

Similarly to Theorem 3, we let  $k = (s + 1)^a k'$  and  $m + sk + 1 = (s + 1)^b m'$ , where  $k'$  and  $m'$  are not divisible by  $(s + 1)$ . As  $s + 1$  is prime,  $(s + 1, k') = 1$ . Therefore, the following result can be derived immediately from Theorems 3 and 13, and Corollary 12.

**Theorem 14** *Suppose  $m \geq (s+1)k$ ,  $s+1$  is prime, and  $m, k, a, b$  are defined as above. Then*

$$\chi(Z, D_{m,k,s}) = \begin{cases} \lceil (m+sk+1)/(s+1) \rceil, & \text{if } a=0 \text{ or } a < b; \\ (m+sk+1)/(s+1) + 1, & \text{if } 0 < b \leq a. \end{cases}$$

Suppose  $k$  is a power of a prime,  $k = p^a$ . If  $p \neq s+1$ , by Theorem 14,  $\chi(Z, D_{m,k,s}) = \lceil (m+sk+1)/(s+1) \rceil$  for all  $m \geq (s+1)k$ . If  $p = s+1$ , that is,  $k = (s+1)^a$ , then the chromatic number of  $G(Z, D_{m,k,s})$  can be completely determined as follows.

**Corollary 15** *Suppose  $m \geq (s+1)k$ ,  $s+1$  is prime,  $k = (s+1)^a$ , and  $m+sk+1 = (s+1)^b m'$ , where  $m'$  is not a multiple of  $s+1$ . Then*

$$\chi(Z, D_{m,k,s}) = \begin{cases} \lceil (m+sk+1)/(s+1) \rceil, & \text{if } b=0 \text{ or } a < b; \\ (m+sk+1)/(s+1) + 1, & \text{if } 0 < b \leq a. \end{cases}$$

**Proof.** By Theorem 14, we only have to show the case as  $b=0$ , which implies  $(k, m+sk+1) = 1$ . Hence by Corollary 10, the prove is complete. Q.E.D.

Note that when  $s+1$  is prime, Theorem 14 determines the value of  $\chi(Z, D_{m,k,s})$  unless  $a > 0$  and  $b=0$ . Thus, for the rest of this section, we shall assume that  $a > 0$  and  $b=0$ , that is,  $k$  is a multiple of  $s+1$  but  $m+sk+1$  is not. Our next result completely settles the case for  $a=1$ .

**Theorem 16** *Suppose  $s+1$  is prime, let  $m, s, k, a, b$  be integers same as defined in Theorem 3. If  $a=1$ , then  $\chi(Z, D_{m,k,s}) = \lceil (m+sk+1)/(s+1) \rceil$  for all  $m \geq (s+1)k$ .*

**Proof.** Let  $r = \lceil (m+sk+1)/(s+1) \rceil \bmod (s+1)$ . We consider two cases.

**Case 1.**  $r=0$ . There exists an integer  $\bar{m} \geq m$  such that  $(\bar{m}+sk+1)/(s+1) = \lceil (m+sk+1)/(s+1) \rceil$ . The distance graph  $G(Z, D_{m,k,s})$  is a subgraph of  $G(Z, D_{\bar{m},k,s})$ , so  $\chi(Z, D_{m,k,s}) \leq \chi(Z, D_{\bar{m},k,s})$ . Let  $\bar{m}+sk+1 = (s+1)^{\bar{b}} \bar{m}'$ , where  $\bar{m}'$  is not divisible

by  $(s+1)$ . Since  $(\bar{m} + sk + 1)/(s+1) \equiv r \equiv 0 \pmod{s+1}$ ,  $\bar{b} \geq 2 > 1 = a$ . Thus by Theorems 2 and 3, we have

$$\lceil (m + sk + 1)/(s+1) \rceil \leq \chi(Z, D_{m,k,s}) \leq \chi(Z, D_{\bar{m},k,s}) = (\bar{m} + sk + 1)/(s+1).$$

Therefore,  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s+1) \rceil$ .

**Case 2.**  $1 \leq r \leq s$ . Since  $s+1$  is a prime, there exists an integer  $1 \leq t \leq s$  such that  $tr \equiv 1 \pmod{s+1}$ . Define a pre-coloring  $c$  of the set  $Z^*$  with  $s+1$  colors as follows. For each integer  $j \in Z^*$ , express  $j$  uniquely in the form  $j = u(s+1) + v$ , where  $0 \leq v \leq s$ . Then let  $c(j) = (ut + v) \pmod{s+1}$ . We shall show that  $c$  satisfies (1) and (2) in Corollary 7 with  $n = \lceil (m + sk + 1)/(s+1) \rceil$ .

Let  $j \in Z^*$ . Assume, contrary to (1) of Corollary 7,  $c(j - hk) = c(j - h'k)$  for some  $0 \leq h < h' \leq s$ . Let  $j - hk = u(s+1) + v$  and  $j - h'k = u'(s+1) + v'$ , then  $ut + v \equiv u't + v' \pmod{s+1}$ . Because  $a = 1$ ,  $(s+1)$  divides  $k$ , which implies  $j - hk \equiv j - h'k \pmod{s+1}$ , so  $v = v'$ . Hence,  $ut - u't \equiv 0 \pmod{s+1}$ . This is impossible because  $(t, s+1) = 1$  and  $0 < u' - u \leq s$ .

Now we show that among any  $m - k + 1$  consecutive integers, there are at most  $\lceil (m - k + 1)/(s+1) \rceil$  vertices of pre-color 0. Similarly to the proof of Theorem 13, we divide the set  $Z^*$  into segments of length  $s+1$  by  $A_0 = \{0, 1, \dots, s\}$ ,  $A_1 = \{s+1, s+2, \dots, 2s+1\}$ ,  $\dots$ ,  $A_i = \{i(s+1), i(s+1)+1, \dots, (i+1)(s+1)-1\}$ ,  $\dots$ . Then each of the segments  $A_i$  contains exactly one vertex of each pre-color. Indeed, it is straightforward to verify that the pre-colors of the segment  $A_i$  are  $\{j, j+1, \dots, s, 0, 1, \dots, j-1\}$ , where  $i \equiv v \pmod{s+1}$ ,  $0 \leq v \leq s$ , and  $j = vt \pmod{s+1}$ .

Let  $Y = \{y, y+1, \dots, y+m-k\}$  be a set of  $m-k+1$  consecutive non-negative integers. Suppose  $y \in A_i$  and  $y+m-k \in A_{i'}$ . If  $|Y \cap A_i| + |Y \cap A_{i'}| \geq s+1$ , then  $Y$  intersects  $\lceil (m-k+1)/(s+1) \rceil$  segments. Hence  $Y$  contains at most  $\lceil (m-k+1)/(s+1) \rceil$  vertices of pre-color 0.

Assume  $|Y \cap A_i| + |Y \cap A_{i'}| < s+1$ , then  $i' - i = \lceil (m - k + 1)/(s+1) \rceil \equiv \lceil (m + sk + 1)/(s+1) \rceil \equiv r \pmod{s+1}$ . Recall that  $tr \equiv 1 \pmod{s+1}$ . Hence, if

$A_i$  is pre-colored by colors  $\{j, j + 1, \dots, s, 0, 1, \dots, j - 1\}$ , then  $A_{i'}$  is pre-colored by colors  $\{j + 1, j + 2, \dots, s, 0, 1, \dots, j\}$ . Since  $|Y \cap A_i| + |Y \cap A_{i'}| < s + 1$ , we conclude that pre-color 0 is used at most once in the set  $(Y \cap A_i) \cup (Y \cap A_{i'})$ . Therefore, at most  $\lceil (m - k + 1)/(s + 1) \rceil$  vertices of  $Y$  have pre-color 0. This completes the proof of Theorem 16. Q.E.D.

In the next result, we write  $m - k + 1$  in the form  $m - k + 1 = u(s + 1)k + vk + p(s + 1) + q$ , where  $u, v, p, q$  are integers such that  $u \geq 0$ ,  $0 \leq v \leq s$ ,  $0 \leq p < k/(s + 1)$ ,  $0 \leq q \leq s$ . It is easy to see that the integers  $u, v, p, q$  are uniquely determined by  $m - k + 1$ .

**Theorem 17** *Suppose  $m \geq (s + 1)k$ ,  $k$  is a multiple of the prime  $s + 1$ , but  $m + sk + 1$  is not. Let  $u, v, p, q$  be integers defined as above. If  $q \leq v + 1$ , then  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s + 1) \rceil$ .*

**Proof.** It suffices to show that  $G(Z, D_{m,k,s})$  is  $\lceil (m + sk + 1)/(s + 1) \rceil$ -colorable. Define a pre-coloring as follows. First, partition the set of  $Z^*$  into blocks recursively in such a way that the first  $k$  vertices are divided into  $k - 1$  blocks with  $k - 2$  single-vertex blocks followed by one block with two vertices. Then repeat the same process to the next  $k$  vertices and so on. Next, pre-color the blocks periodically with pre-colors  $\{0, 1, 2, \dots, s\}$ , that is, every vertex in the first block is pre-colored by 0 and so on. It is enough to show that the pre-coloring satisfies (1) and (2) of Corollary 7, with  $n = \lceil (m + sk + 1)/(s + 1) \rceil$ .

First we prove that for any  $j \geq sk$ , the  $s + 1$  vertices  $j, j - k, \dots, j - sk$  receive distinct pre-colors. Suppose  $0 \leq t < t' \leq s$ . Let the pre-colors of  $j - t'k$  and  $j - tk$  be  $x$  and  $y$ , respectively. Because  $s + 1$  divides  $k$ , and  $s + 1$  is prime, we have  $(s + 1, k - 1) = 1$ . As  $(j - tk) - (j - t'k) = (t' - t)k$  and any consecutive  $k$  vertices are divided into  $k - 1$  blocks, so  $y \equiv x + (t' - t)(k - 1) \pmod{s + 1}$ . Hence, we conclude that  $x \neq y$ , since  $1 \leq t' - t < s + 1$  and  $(s + 1, k - 1) = 1$ .

Next we prove that among any  $m - k + 1$  consecutive vertices, there are at most  $\lceil (m - k + 1)/(s + 1) \rceil$  vertices with pre-color 0. Given a set  $Y$  of  $m - k + 1$  consecutive non-negative integers, we may assume that the first two vertices of  $Y$  have pre-color 0. Among the first  $u(s + 1)k$  vertices of  $Y$ , exactly  $uk$  of them have pre-color 0, because any consecutive  $(s + 1)k$  vertices are evenly pre-colored, *i.e.*, there are exactly  $k$  vertices of each pre-color.

The assumption that  $m + sk + 1$  is not a multiple of  $s + 1$  implies that  $m - k + 1$  is not a multiple of  $s + 1$ . Because  $k$  is a multiple of  $s + 1$  while  $m - k + 1$  is not,  $p(s + 1) + q \geq 1$ . If  $p(s + 1) + q \geq 2$ , then among the remaining  $vk + p(s + 1) + q$  vertices of  $Y$ , there are  $v + 1$  blocks of size 2. If we remove one vertex from each of these blocks of size 2, then the remaining  $vk + p(s + 1) + q - v - 1$  vertices of  $Y$  are *almost* evenly pre-colored, that is, the numbers of vertices with same pre-colors differ by at most one. Hence at most  $\lceil (vk + p(s + 1) + q - v - 1)/(s + 1) \rceil$  of them have pre-color 0. On the other hand, among the removed vertices, exactly one vertex has pre-color 0. Therefore, the total number of vertices of pre-color 0 is at most  $uk + 1 + \lceil (vk + p(s + 1) + q - v - 1)/(s + 1) \rceil = \lceil (m - k + 1)/(s + 1) \rceil$ . Note that the last equality is due to the assumption that  $q \leq v + 1$ .

Finally, we assume  $p(s + 1) + q = 1$ . Then it is straightforward to verify that either  $v = 0$ , or the pre-color of the last vertex is not 0. Consider the remaining  $vk + p(s + 1) + q = vk + 1$  vertices of  $Y$ . If  $v = 0$ , then there is one vertex of pre-color 0. If the pre-color of the last vertex is not 0, then among the remaining  $vk + 1$  vertices of  $Y$ , there are  $v$  blocks of size 2. If we remove one vertex from each of these blocks of size 2, then the remaining  $vk - v$  vertices of  $Y$  are almost evenly pre-colored, so at most  $\lceil (vk - v)/(s + 1) \rceil$  of them have pre-color 0. On the other hand, among the vertices taken away, only one has pre-color 0. Hence, there are at most  $1 + \lceil (vk - v)/(s + 1) \rceil = \lceil (vk + 1)/(s + 1) \rceil$  (because  $v \leq s$ ) vertices of pre-color 0 in the remaining  $vk + 1$  vertices of  $Y$ . Therefore, we conclude that  $Y$  has at most

$uk + \lceil (vk + 1)/(s + 1) \rceil = \lceil (m - k + 1)/(s + 1) \rceil$  vertices with pre-color 0. This completes the proof. Q.E.D.

**Corollary 18** *Suppose  $m \geq (s + 1)k$ ,  $k$  is a multiple of the prime  $s + 1$ , but  $m + sk + 1$  is not. Let  $u, v, p, q$  be the same as defined in Theorem 17. If  $v \geq s - 1$ , or  $q \leq 1$ , then  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s + 1) \rceil$ .*

Note that when  $s = 1$ , then  $v \geq s - 1$  is always true, hence we have the following corollary which was proved in [2]:

**Corollary 19** *Suppose  $s = 1$ ,  $m \geq 2k$ ,  $k = 2^a k'$  and  $m + k + 1 = 2^b m'$ , where  $k'$  and  $m'$  are odd. Then*

$$\chi(Z, D_{m,k,1}) = \begin{cases} \lceil (m + k + 1)/2 \rceil, & \text{if } b = 0 \text{ or } a < b; \\ ((m + k + 1)/2) + 1, & \text{if } 0 < b \leq a. \end{cases}$$

**Proof.** The case as  $b = 0$  follows from Corollary 18; and the case as  $b > 0$  follows from Theorem 14. Q.E.D.

Recall that  $k = (s + 1)^a k'$  where  $a \geq 1$  and  $k'$  is not divisible by  $s + 1$ , and  $m - k + 1$  is not divisible by  $s + 1$ . In order to introduce the next result, we need the following definitions and notations. For any factor  $x$  of  $k'$ , define:

$$\begin{aligned} q(x) &:= \lceil (m - k + 1)/((s + 1)^a x) \rceil \bmod (s + 1); \\ m(t, x) &:= \max\{t(q(x) - 1) \bmod (s + 1), tq(x) \bmod (s + 1)\}, 1 \leq t \leq s; \\ f(x) &:= \min\{m(t, x) : 1 \leq t \leq s\}. \end{aligned}$$

Finally, define  $f := \min\{f(x) : x \text{ is a factor of } k'\}$ .

Note that for given  $m, k$  and  $s$ , the integer  $f$  in the above is uniquely determined. Similarly as in Theorem 17, we let  $q = (m - k + 1) \bmod (s + 1)$ .

**Theorem 20** *Given  $m, k$  and  $s$  where  $m \geq (s + 1)k$  and  $s + 1$  is a prime, let  $f, q$  be defined as above. If  $f + q \leq s + 1$ , then  $\chi(Z, D_{m,k,s}) = \lceil \chi_f(Z, D_{m,k,s}) \rceil = \lceil (m + sk + 1)/(s + 1) \rceil$ .*

**Proof.** Suppose  $f = f(x) = m(t, x)$  for some factor  $x$  of  $k'$  and some  $1 \leq t \leq s$ . Express any integer  $j \in Z^*$  in the following form:

$$j = u(s+1)^a x + v(s+1) + w,$$

where  $u \geq 0$ ,  $0 \leq v < (s+1)^{a-1}x$  and  $0 \leq w \leq s$ .

It is easy to see that for each  $j$ , the integers  $u, v, w$  in the form above are uniquely determined by  $j$ . Define a pre-coloring  $c$  using the  $s+1$  pre-colors  $\{0, 1, \dots, s\}$  by  $c(j) = (ut + w) \bmod (s+1)$ . In order to prove  $G(Z, D_{m,k,s})$  is  $\lceil (m+sk+1)/(s+1) \rceil$ -colorable, it suffices to show that  $c$  satisfies (1) and (2) of Corollary 7, with  $n = \lceil (m+sk+1)/(s+1) \rceil$ .

First, let  $j$  be any non-negative integer, we shall show that  $c(j), c(j-k), c(j-2k), \dots, c(j-sk)$  are all distinct. Let  $0 \leq p' < p \leq s$ . If  $j - pk = u(s+1)^a x + v(s+1) + w$ , then

$$\begin{aligned} j - p'k &= u(s+1)^a x + v(s+1) + w + (p - p')k \\ &= u(s+1)^a x + v(s+1) + w + (p - p')(s+1)^a k' \\ &= u'(s+1)^a x + v(s+1) + w. \end{aligned}$$

Because  $(s+1, k') = (p - p', s+1) = 1$ , one has  $(u' - u, s+1) = 1$ . Assume  $c(j - pk) = c(j - p'k)$ , then  $ut + w \equiv u't + w \pmod{s+1}$ . Hence  $t(u' - u) \equiv 0 \pmod{s+1}$ , which is impossible, since  $s+1$  is prime and  $(t, s+1) = (u' - u, s+1) = 1$ . This proves that  $c$  satisfies (1) of Corollary 7.

Next, we prove that among any  $m - k + 1$  consecutive integers, there are at most  $\lceil (m - k + 1)/(s+1) \rceil$  vertices with pre-color 0. Divide the vertex set  $Z^*$  evenly into segments of length  $s+1$  by  $A_0 = \{0, 1, 2, \dots, s\}, A_1 = \{s+1, s+2, \dots, 2s+1\}, \dots, A_i = \{i(s+1), i(s+1)+1, \dots, (i+1)(s+1)-1\}, \dots$ . Then each of the segments  $A_i$  contains exactly one vertex of each pre-color. Indeed, the pre-colors of the segment  $A_i$  are  $\{j, j+1, \dots, s, 0, 1, \dots, j-1\}$ , where  $j = ut \bmod (s+1)$ , and  $u$  is the unique integer such that  $i = u(s+1)^{a-1}x + v$ ,  $0 \leq v < (s+1)^{a-1}x$ .

Let  $Y$  be a set of  $m - k + 1$  consecutive integers,  $Y = \{y, y+1, \dots, y+m-k\}$ .

Suppose  $y \in A_i$  and  $y + m - k \in A_{i'}$ . If  $|Y \cap A_i| + |Y \cap A_{i'}| \geq s + 1$ , then  $Y$  has at most  $\lceil (m - k + 1)/(s + 1) \rceil$  vertices with pre-color 0 (cf. proof of Theorem 16).

Now we assume that  $|Y \cap A_i| + |Y \cap A_{i'}| < s + 1$ , then  $|Y \cap A_i| + |Y \cap A_{i'}| = q$ . Suppose  $i = u(s + 1)^{a-1}x + v$  and  $i' = u'(s + 1)^{a-1}x + v'$ , where  $0 \leq v, v' < (s + 1)^{a-1}x$ . Then by the definition of  $q(x)$ , either  $u' - u = q(x)$  or  $u' - u = q(x) - 1$ . Suppose  $\alpha = q(x)t \bmod (s + 1)$  and  $\beta = (q(x) - 1)t \bmod (s + 1)$ . Then by the choice of  $x$  and  $t$ , one has  $\alpha, \beta \leq f$ .

Suppose the pre-colors of  $A_i$  are  $\{j, j + 1, \dots, s, 0, 1, \dots, j - 1\}$ . Then the pre-colors of  $A_{i'}$  are either  $\{j + \alpha, j + \alpha + 1, \dots, s, 0, 1, \dots, j + \alpha - 1\}$ , if  $u' - u = q(x)$ ; or  $\{j + \beta, j + \beta + 1, \dots, s, 0, 1, \dots, j + \beta - 1\}$ , if  $u' - u = q(x) - 1$ .

Any other segment different from  $A_i$  and  $A_{i'}$  is either disjoint from  $Y$  or contained in  $Y$ . As each segment contains exactly one vertex of each color, to prove that  $Y$  has at most  $\lceil (m - k + 1)/(s + 1) \rceil$  vertices with pre-color 0, it suffices to show that the pre-color 0 is used at most once in the union  $(Y \cap A_i) \cup (Y \cap A_{i'})$ . Assume that 0 is used in both  $Y \cap A_i$  and  $Y \cap A_{i'}$ . Without loss of generality, we may assume that the pre-colors of  $A_{i'}$  are  $\{j + \alpha, j + \alpha + 1, \dots, s, 0, 1, \dots, j + \alpha - 1\}$ . Then one has  $|Y \cap A_i| \geq j$  and  $|Y \cap A_{i'}| \geq s + 1 - (j + \alpha - 1)$ . It follows that  $q = |(Y \cap A_i) \cup (Y \cap A_{i'})| \geq s + 2 - \alpha$ , contrary to the assumption that  $\alpha + q \leq f + q \leq s + 1$ . Therefore  $c$  satisfies (2) of Corollary 7, with  $n = \lceil (m + sk + 1)/(s + 1) \rceil$ . This completes the proof of Theorem 20. Q.E.D.

**Corollary 21** *If  $m \geq (s + 1)k$ ,  $s + 1$  is prime, and there is a factor  $x$  of  $k'$  such that  $q(x) \leq 1$ , then  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s + 1) \rceil$ . In particular, if  $\lceil (m - k + 1)/k \rceil \bmod (s + 1) \leq 1$ , then  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s + 1) \rceil$ .*

**Proof.** According to definition, if  $q(x) = 1$ , then  $m(1, x) = 1$ ; if  $q(x) = 0$ , then  $m(t, x) = 1$  for some  $t$  such that  $ts \equiv 1 \pmod{s + 1}$ . (Such a  $t$  exists, because

$(s, s + 1) = 1$ .) In any of the two cases,  $f = 1$ , so  $f + q \leq s + 1$ . Therefore,  $\chi(Z, D_{m,k,s}) = \lceil (m + sk + 1)/(s + 1) \rceil$  by Theorem 20. Q.E.D.

Applying Theorem 14 and Corollaries 18 and 21, we are able to completely settle the case  $s = 2$ .

**Corollary 22** *Suppose  $s = 2$ ,  $m \geq 3k$ ,  $k = 3^a k'$  and  $m + 2k + 1 = 3^b m'$ , where  $k'$  and  $m'$  are not multiples of 3. Then*

$$\chi(Z, D_{m,k,2}) = \begin{cases} \lceil (m + 2k + 1)/3 \rceil, & \text{if } b = 0 \text{ or } a < b; \\ (m + 2k + 1)/3 + 1, & \text{if } 0 < b \leq a. \end{cases}$$

**Proof.** According to Theorem 14, we only have to show the case as  $b = 0$ . Suppose  $m - k + 1 = u(s + 1)k + vk + p(s + 1) + q$ . If  $v \neq 0$ , then the conclusion follows from Corollary 18. If  $v = 0$ , then the conclusion follows from Corollary 21. (Because  $\lceil (m - k + 1)/k \rceil \bmod (s + 1) \leq 1$ .) Q.E.D.

**Remarks.** New results related to this topic have been obtained since the submission of this paper. In [5], it was proved that  $\chi(G(Z, D_{m,k,s})) \leq \lceil (m + sk + 1)/(s + 1) \rceil + 1$  for all  $m \geq (s + 1)k$ . Then in [11], the chromatic numbers of all the graphs  $G(Z, D_{m,k,s})$  are completely determined. The circular chromatic number of the class of distance graphs  $G(Z, D_{m,k,s})$  was studied in [1, 11, 19], and the value of  $\chi_c(Z, D_{m,k,s})$  has been completely determined in [19]. (The circular chromatic number  $\chi_c(G)$  of a graph  $G$  is a refinement of  $\chi(G)$ , and  $\chi(G) = \lceil \chi_c(G) \rceil$  for any graph  $G$ . For a survey of research concerning circular chromatic number of graphs, see [20].)

**Acknowledgments.** The authors are grateful to the referees for many helpful suggestions.

## References

- [1] G. J. Chang, L. Huang and X. Zhu, *The circular chromatic numbers and the fractional chromatic numbers of distance graphs*, Europ. J. of Comb., to appear.

- [2] G. J. Chang, D. D.-F. Liu and X. Zhu, *Distance graphs and T-coloring*, J. of Comb. Theory, Series B, to appear.
- [3] J. J. Chen, G. J. Chang and K. C. Huang, *Integral distance graphs*, J. Graph Theory 25 (1997) 287-294.
- [4] W. Deuber and X. Zhu, *The chromatic number of distance graphs*, Disc. Math. 165/166 (1997) 195-204.
- [5] W. Deuber and X. Zhu, *Chromatic numbers of distance graphs with distance sets missing multiples*, manuscript, 1997.
- [6] R. B. Eggleton, P. Erdős and D. K. Skilton, *Coloring the real line*, J. Comb. Theory, Series B 39 (1985) 86-100.
- [7] R. B. Eggleton, P. Erdős and D. K. Skilton, *Research problem 77*, Disc. Math. 58 (1986) 323.
- [8] R. B. Eggleton, P. Erdős and D. K. Skilton, *Update information on research problem 77*, Disc. Math. 69 (1988) 105.
- [9] R. B. Eggleton, P. Erdős and D. K. Skilton, *Coloring prime distance graphs*, Graphs and Comb. 32 (1990) 17-32.
- [10] J. R. Griggs and D. D.-F. Liu, *The channel assignment problem for mutually adjacent sites*, J. Comb. Theory, Series A, 68 (1994) 169-183.
- [11] L. Huang and G. J. Chang, *Circular chromatic number of distance graphs with distance set missing multiples*, manuscript, 1998.
- [12] A. Kemnitz and H. Kolberg, *Coloring of integer distance graphs*, Disc. Math., to appear (1998).

- [13] D. D.-F. Liu, *T-coloring and chromatic number of distance graphs*, Ars. Comb., to appear.
- [14] E. R. Scheinerman and D. H. Ullman, *Fractional Graph Theory*, Wiley-Interscience Series in Discrete Mathematics and Optimization, 1997.
- [15] M. Voigt, *Coloring of distance graphs*, Ars Comb., to appear.
- [16] M. Voigt and H. Walther, *Chromatic number of prime distance graphs*, Disc. Appl. Math. 51 (1994) 197-209.
- [17] X. Zhu, *Distance graphs on the real line*, manuscript, 1996.
- [18] X. Zhu, *Pattern-periodic coloring of distance graphs*, J. of Comb. Theory, Series B, 73 (1998), 195-206.
- [19] X. Zhu, *The circular chromatic number of a class of distance graphs*, manuscript, 1998.
- [20] X. Zhu, *Circular chromatic number: A survey*, manuscript, 1997.